



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
28.02.2001 Bulletin 2001/09

(21) Application number: **92910933.8**

(22) Date of filing: **18.03.1992**

(51) Int Cl.7: **C12N 5/00, C12N 15/00,
C12N 15/63, C12N 15/74,
C12N 15/80, C12N 1/15,
C12N 1/21, C12P 7/06,
C12P 7/08, C07H 15/12**

(86) International application number:
PCT/US92/01807

(87) International publication number:
WO 92/16615 (01.10.1992 Gazette 1992/25)

(54) **ETHANOL PRODUCTION BY RECOMBINANT HOSTS**

ETHANOLHERSTELLUNG DURCH REKOMBINANTE WIRTE

PRODUCTION D'ETHANOL PAR DES MICROORGANISMES HOTES DE RECOMBINAISON

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU MC NL SE

(30) Priority: **18.03.1991 US 670821**
06.03.1992 US 846344

(43) Date of publication of application:
05.01.1994 Bulletin 1994/01

(73) Proprietors:
• **UNIVERSITY OF FLORIDA**
Gainesville, Florida 32611-2037 (US)
• **BIOENERGY INTERNATIONAL, L.C.**
Gainsville, FL 32606-1657 (US)

(72) Inventors:
• **INGRAM, Lonnie, O.**
Gainesville, FL 32606 (US)
• **BEALL, David, S.**
Gainesville, FL 32601 (US)
• **BURCHHARDT, Gerhard, F., H.**
Gainesville, FL 32605 (US)
• **GUIMARAES, Walter, V.**
Gainesville, FL 32611 (US)
• **OHTA, Kazuyoshi**
Gainesville, FL 32611 (US)
• **WOOD, Brent, E.**
Gainesville, FL 32601 (US)
• **SHANMUGAM, Keelnatham, T.**
Gainesville, FL 32605 (US)
• **FOWLER, David, A.**
Gainesville, FL 32608 (US)

• **BEN-BASSAT, Arie**
Gainesville, FL 32606 (US)

(74) Representative: **Price, Vincent Andrew et al**
FRY HEATH & SPENCE
The Old College
53 High Street
Horley Surrey RH6 7BN (GB)

(56) References cited:
WO-A-90/02193 WO-A-92/10561
US-A- 4 493 893 US-A- 4 612 287
US-A- 4 839 286

- **Appl. Environ. Microbiol., Volume 53, No. 10,**
issued October 1987, INGRAM et al., "Genetic
Engineering of Ethanol Production in
Escherichia coli", pages 2420-2425, see entire
document.
- **Appl. Environ. Microbiol., Volume 53, No. 9,**
issued September 1987, TOLAN et al.,
"Fermentation of D-Xylose and L-Arabinose to
Ethanol by Erwinia crysanthemii", pages
2033-2038, see entire document.
- **Nucleic Acids Res., Volume 15, No. 4, issued 25**
February 1987, NEALE et al., "Nucleotide
Sequence of the Pyruvate Decarboxylase Gene
from Zymomonas mobilis", pages 1753-1761,
see entire document.

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- Arch. Microbiol., Volume 144, issued 1986, BRAU et al., "Cloning and Expression of the Structural Gene for Pyruvate Decarboxylase of *Zymomonas mobilis* in *E. coli*", pages 296-301, see entire document.
- J. Bacteriol., Volume 169, No. 6, issued June 1987, CONWAY et al., "Cloning and Sequencing of the Alcohol Dehydrogenase II Gene from *Zymomonas mobilis*", pages 2591-2597, see entire document.
- Appl. Environ. Microbiol., Volume 54, No. 2, issued February 1988, CURRY et al., "Expression and Secretion of Cellulomonas fimi Exoglucanase in *Saccharomyces cerevisiae*", pages 476-484, see entire document.
- Appl. Environ. Microbiol., Volume 53, No. 2, issued February 1987, CONWAY et al., "Expression Vector for *Zymomonas mobilis*", pages 235-241, see entire document.
- Appl. Environ. Microbiol., Volume 53, No. 9, issued September 1987, TOLAN et al., "Fermentation of D-Xylose to Ethanol by Genetically Modified *Klebsiella planticola*", pages 2039-2044, see entire document.
- Bio/Technology, Volume 4, issued October 1986, JOLIFF et al., "Isolation, Crystallization and Properties of a New Cellulase of *Clostridium thermocellum* overproduced in *Escherichia coli*", pages 896-900, see entire document.
- Bio/Technology, Volume 1, issued September 1983, CORNET et al., "Characterization of Two cel (Cellulose Degradation) Genes of *Clostridium thermocellum* Coding for Endoglucanases", pages 589-594, see entire document.
- J. Biotechnol., Volume 12, issued 1989, AL-ZAAG, "Molecular Cloning of Cellulose and Other beta-glucosidase Determinants from *Klebsiella ocytoca*", pages 79-86, see entire document.
- Appl. Environ. Microbiol., Volume 55, No. 1, issued January 1989, TAILLIEZ et al., "Cellulose Fermentation by an Asporogenous Mutant and an Ethanol-Tolerant Mutant of *Clostridium thermocellum*", pages 203-206, see entire document.
- Biochimie, Volume 68, issued 1986, PETRE et al., "Purification and Properties of the Endoglucanase C of *Clostridium thermocellum* Produced in *Escherichia coli*", pages 687-695, see entire document.
- J. Bacteriol., Volume 170, No. 10, issued October 1988, GREPINET et al., "Nucleotide Sequence and Deletion Analysis of the Xylanase Gene (xynZ) of *Clostridium thermocellum*", pages 4582-4588, see entire document.
- J. Bacteriol., Volume 170, No. 10, issued October 1988, GREPINET et al., "Purification of *Clostridium thermocellum* Xylanase Z Expressed in *Escherichia coli* and Identification of the Corresponding Product in the Culture Medium of *C. thermocellum*", see entire document.
- FEMS Microbiol. Lett., Volume 29, issued 1985, MILLET et al., "Cloning of Ten Distinct DNA Fragments of *Clostridium thermocellum* Coding for Cellulases", pages 145-149, see entire document.

Description

Background of the Invention

[0001] During glycolysis, cells convert simple sugars like glucose into pyruvic acid, with a net production of ATP and NADH. In the absence of a functioning electron transport system for oxidative phosphorylation, at least 95% of the pyruvic acid is consumed in short pathways which regenerate NAD⁺, an obligate requirement for continued glycolysis and ATP production. The waste products of these NAD⁺ regeneration systems are commonly referred to as fermentation products.

[0002] Microorganisms are particularly diverse in the array of fermentation products which are produced by different genera. These products include organic acids, such as lactate, acetate, succinate and butyrate, as well as neutral products such as ethanol, butanol, acetone and butanediol. Indeed, the diversity of fermentation products from bacteria has led to their use as a primary determinant in taxonomy. See, for example, *BERGEY'S MANUAL OF SYSTEMATIC BACTERIOLOGY*, Williams & Wilkins Co., Baltimore (1984) (hereafter "Bergey's Manual").

[0003] End products of fermentation share several fundamental features. They are relatively nontoxic under the conditions in which they are initially produced but become more toxic upon accumulation. They are more reduced than pyruvate because their immediate precursors have served as terminal electron acceptors during glycolysis. The microbial production of these fermentation products forms the basis for our oldest and most economically successful applications of biotechnology and includes dairy products, meats, beverages and fuels. In recent years, many advances have been made in the field of biotechnology as a result of new technologies which enable researchers to selectively alter the genetic makeup of some microorganisms.

[0004] The bacterium *Escherichia coli* is an important vehicle for the cloning and modification of genes for biotechnology, and is one of the most important hosts for the production of recombinant products. In recent years, the range of hosts used for recombinant DNA research has been extended to include a variety of bacteria, yeasts, fungi, and eukaryotic cells. The invention described here relates to the use of recombinant DNA technology to elicit the production of specific useful products by a modified host.

[0005] The DNA used to modify the host of the present invention can be isolated from *Zymomonas mobilis*, a bacterium which commonly is found in plant saps and in honey. *Z. mobilis* has long served as an inoculum for palm wines and for the fermentation of *Agave* sap to produce *pulque*, an alcohol-containing Mexican beverage. The microbe also is used in the production of fuel ethanol, and reportedly is capable of ethanol-production rates which are substantially higher than that of yeasts.

[0006] Although *Z. mobilis* is nutritionally simple and capable of synthesizing amino acids, nucleotides and vitamins, the range of sugars metabolized by this organism is very limited and normally consists of glucose, fructose and sucrose. Substrate level phosphorylation from the fermentation of these sugars is the sole source of energy for biosynthesis and homeostasis. *Zymomonas mobilis* is incapable of growth even in rich medium such as nutrient broth without a fermentable sugar.

[0007] *Z. mobilis* is an obligatively fermentative bacterium which lacks a functional system for oxidative phosphorylation. Like the yeast *Saccharomyces cerevisiae*, *Z. mobilis* produces ethanol and carbon dioxide as principal fermentation products. *Z. mobilis* produces ethanol by a short pathway which requires only two enzymatic activities: pyruvate decarboxylase and alcohol dehydrogenase. Pyruvate decarboxylase is the key enzyme in this pathway which diverts the flow of pyruvate to ethanol. Pyruvate decarboxylase catalyzes the nonoxidative decarboxylation of pyruvate to produce acetaldehyde and carbon dioxide. Two alcohol dehydrogenase isozymes are present in this organism and catalyze the reduction of acetaldehyde to ethanol during fermentation, accompanied by the oxidation of NADH to NAD⁺. Although bacterial alcohol dehydrogenases are common in many organisms, few bacteria have pyruvate decarboxylase. Attempts to modify *Z. mobilis* to enhance its commercial utility as an ethanol producer have met with very limited success.

[0008] Most fuel ethanol is currently produced from hexose sugars in corn starch or cane syrup utilizing *S. cerevisiae* or *Z. mobilis*. But such sugars are relatively expensive sources of biomass sugars and have competing value as foods. Starch and sugars represent only a fraction of the total carbohydrates in plants. The dominant forms of plant carbohydrate in stems, leaves, hulls, husks, cobs and the like are the structural wall polymers, cellulose and hemicellulose. Hydrolysis of these polymers releases a mixture of neutral sugars which include glucose, xylose, mannose, galactose, and arabinose. No single organism has been found in nature which can rapidly and efficiently metabolize all of these sugars into ethanol or any other single product of value.

[0009] The genes coding for alcohol dehydrogenase II and pyruvate decarboxylase in *Z. mobilis* have been separately cloned, characterized, and expressed in *E. coli*. See Brau & Sahm [1986a] *Arch. Microbiol.* 144: 296-301, [1986b] *Arch. Microbiol.* 146: 105-10; Conway *et al.* [1987a] *J. Bacteriol.* 169: 949-54; Conway *et al.* [1987b] *J. Bacteriol.* 169: 2591-97; Neale *et al.* [1987] *Nucleic Acid. Res.* 15: 1753-61; Ingram & Conway [1988] *Appl. Environ. Microbiol.* 54: 397-404; Ingram *et al.* [1987] *Appl. Environ. Microbiol.* 53: 2420-25.

[0010] Brau and Sahm [1986a], *supra*, first demonstrated that ethanol production could be increased in recombinant *E. coli* by the over-expression of *Z. mobilis* pyruvate decarboxylase although very low ethanol concentrations were produced. Subsequent studies extended this work by using two other enteric bacteria, *Erwinia chrysanthemi* and *Klebsiella planticola*, and thereby achieved higher levels of ethanol from hexoses, pentoses, and sugar mixtures. See Tolan & Finn [1987] *Appl. Environ. Microbiol.* 53: 2033-38, 2039-44.

[0011] When a feedstock of simple sugars is available, therefore, the aforementioned microbes generally are useful. Nevertheless, the majority of the world's cheap, renewable source of biomass is not found as monosaccharides but rather in the form of lignocellulose, which is primarily a mixture of cellulose, hemicellulose, and lignin. Cellulose is a homopolymer of glucose, while hemicellulose is a more complex heteropolymer comprised not only of xylose, which is its primary constituent, but also of significant amounts of arabinose, mannose, glucose and galactose. It has been estimated that microbial conversion of the sugar residues present in waste paper and yard trash from U.S. landfills could provide over ten billion gallons of ethanol. While microorganisms such as those discussed above can ferment efficiently the monomeric sugars which make up the cellulosic and hemicellulosic polymers present in lignocellulose, the development of improved methods for the saccharification of lignocellulose remains a major research goal.

[0012] Current methods of saccharifying lignocellulose include acidic and enzymatic hydrolyses. Acid hydrolysis usually requires heat and presents several drawbacks, however, including the use of energy, the production of acidic waste, and the formation of toxic compounds which can hinder subsequent microbial fermentations. Enzymatic hydrolysis thus presents a desirable alternative. For example, enzymes can be added directly to the medium containing the lignocellulosic material.

[0013] Genetic-engineering approaches for the addition of saccharifying traits to microorganisms for the production of ethanol or lactic acid have been directed at the secretion of high enzyme levels into the medium. That is, the art has concerned itself with modifying microorganisms already possessing the requisite proteins for transporting cellularly-produced enzymes into the fermentation medium, where those enzymes can then act on the polysaccharide substrate to yield mono- and oligosaccharides. This approach has been taken because the art has perceived difficulty in successfully modifying organisms lacking the requisite ability to transport such proteins.

Summary of the Invention

[0014] It is therefore an object of the present invention to provide microorganisms that are capable of effectively diverting pyruvate to ethanol during growth under both aerobic and anaerobic conditions.

[0015] It is also an object of the present invention to increase the growth of recombinant hosts and to reduce accumulation of undesirable metabolic products in the growth medium of these hosts.

[0016] It is yet another object of the present invention to provide recombinant hosts that produce sufficient intracellular levels of polysaccharase enzyme(s) to effect the breakdown of glucose or xylose oligomers to products which the same hosts can ferment to ethanol.

[0017] In achieving these and other objects, there is provided, in accordance with the first major domain of the present invention, a recombinant host, other than *Escherichia coli*, comprising heterologous DNA coding for production of alcohol dehydrogenase and pyruvate decarboxylase and polysaccharase or a combination of different polysaccharases in the host. The host expresses the DNA at a sufficient functional level so as to facilitate the production of ethanol as a primary fermentation product of the host.

[0018] One advantageous embodiment employs the *Z. mobilis* genes coding for alcohol dehydrogenase and pyruvate decarboxylase in the recombinant hosts. Other aspects of the first domain include plasmids comprising genes coding for alcohol dehydrogenase, pyruvate decarboxylase and polysaccharase which plasmids are useful for providing the inventive recombinant hosts.

[0019] In the second major domain of this invention, the recombinant host, in addition to comprising the DNA coding for alcohol dehydrogenase, pyruvate decarboxylase and polysaccharase, further comprises DNA coding for proteins which enable the host to transport and metabolize an oligosaccharide. The host expresses the DNA at a level such that the host produces ethanol as a primary fermentation product from the metabolism of the oligosaccharide. Preferred hosts of this domain include enteric bacteria such as *Erwinia* and *Klebsiella*. Other preferred hosts will metabolize a di- and/or trisaccharide comprising C₅ and/or C₆ sugar monomers such as glucose, xylose and maltose.

[0020] The production of a polysaccharase by the host, and the subsequent release of that polysaccharase into the medium, will reduce the amount of commercial enzyme necessary to degrade the feedstock into fermentable monosaccharides and oligosaccharides.

[0021] The polysaccharase DNA can be native to the host, although more often the DNA will be foreign, i.e., heterologous. Advantageous polysaccharases include cellulolytic, xylanolytic and starch-degrading enzymes. The polysaccharase can be at least partially secreted by the host, or it can be accumulated substantially intracellularly for subsequent release. Advantageously, intracellularly-accumulated enzymes which are thermostable, can be released when desired by heat-induced lysis. Combinations of enzymes can be encoded by the heterologous DNA, some of which

are secreted, and some of which are accumulated.

[0022] Other modifications can be made to enhance the ethanol production of the foregoing hosts. For example, the host can further comprise an additional heterologous DNA segment, the expression product of which is a protein involved in the transport of mono- and/or oligosaccharides into the recombinant host. Likewise, additional genes from the glycolytic pathway can be incorporated into the host. In such ways, an enhanced rate of ethanol production can be achieved.

[0023] Also, methods for producing ethanol using all of the foregoing hosts are provided. Further, methods are provided for treating an oligosaccharide feedstock to convert the oligosaccharide into ethanol, and into simpler oligosaccharides and/or saccharide monomers. Yet another aspect provides a method for reducing the accumulation of acidic metabolic products in a growth medium by employing the inventive transformed hosts to produce ethanol in the medium. Another aspect of this invention includes process designs for fermenting oligosaccharide-containing biomass to ethanol.

[0024] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

Brief Description of the Drawings

[0025] Figure 1 illustrates the construction of pLOI295 containing genes encoding pyruvate decarboxylase and alcohol dehydrogenase II from *Z. mobilis*. Abbreviations: R1, *EcoRI*; H, *HindIII*; B, *BamHI*; t, terminator; *adh*, *Z. mobilis* alcohol dehydrogenase II; *pdc*, *Z. mobilis* pyruvate decarboxylase; *cat*, chloramphenicol acyltransferase; Ap, β -lactamase; Zm Pro frags, fragments of *Z. mobilis* DNA which exhibit promoter activity; *ColE1*, replication origin derived from pBR322; *oriV*, replication origin derived from RSF1010; *P_{lac}*, lac promoter; *P_{Zm}*, promoter from *Z. mobilis*; *P_γ*, cryptic promoter on vector; Kb, kilobases.

[0026] Figure 2 illustrates several plasmids containing the genes encoding pyruvate decarboxylase and alcohol dehydrogenase II from *Z. mobilis*.

[0027] Figure 3 illustrates growth and acid production by strain TC4 and recombinants containing plasmids encoding ethanologenic enzymes.

[0028] Figure 4 illustrates the relationship between pyruvate decarboxylase activity in recombinants and the extent of growth. Cell mass after 24 hours of growth is expressed as the optical density at 550 nm (O.D. 550).

[0029] Figure 5 illustrates the construction of pLOI510.

[0030] Figure 6 illustrates the concentration of ethanol produced from fermentation of cellobiose by ethanologenic recombinants of *Erwinia* and of *Klebsiella*.

[0031] Figures 7A, 7B and 7C illustrate the construction of recombinant plasmids containing *xynZ* and *xyfB*.

[0032] Figures 8A and 8B comprise thin layer chromatographic analysis of xylan hydrolysis using *E. coli* strain KO11 (pLOI2003) incubated at 45°C (Fig. 8A) or 60°C (Fig. 8B) in Luria broth (pH 6.0) containing 4% birchwood xylan.

[0033] Figures 9A and 9B illustrate the utilization of xylooligosaccharides for growth by recombinant *E. coli* KO11 (Figure 9A) and *K. oxytoca* M5A1 (Figure 9B).

[0034] Figures 10A and 10B illustrate the conversion of xylose and xylan hydrolysate to ethanol by recombinant *K. oxytoca* M5A1. Figure 10A illustrates the growth and Figure 10B illustrates the ethanol production.

[0035] Figures 11A and 11B illustrate the ethanol production by recombinant strains of *K. oxytoca* M5A1. Figure 11A illustrates the ethanol production from glucose (100 g/liter) by strain M5A1(pLOI555), strain P2 containing integrated *pet* genes, and strain B1 containing integrated *pet* genes. Figure 11B illustrates the ethanol production from cellobiose (100 g/liter) fermentation by strain P2.

[0036] Figure 12 illustrates the effect of commercial cellulase additions on ethanol yield from 50 g SOLKA FLOC SW40/liter after 120 hours.

[0037] Figures 13A, B and C comprise the thin layer chromatographic analyses of cellulose hydrolysis and fermentation by P2(pCT603T) of acid swollen cellulose (Fig. 13A), base swollen cellulose (Fig. 13B), and crystalline cellulose (Fig. 13C).

[0038] Figure 14 illustrates the enhancing effect on ethanol production of pretreatment with the *celD* gene product and using 1% cellulase, which provides an ethanol yield equivalent to that of 5% CYTOLASE.

[0039] Figure 15 illustrates the construction of plasmids pLOI140 and pLOI141, which comprise the α -amylase gene from *Bacillus stearothermophilus* and the pullulanase gene from *Thermoanaerobium brockii*.

[0040] Figures 16A and B illustrate ethanol production and cell mass, respectively, for the fermentation of glucose and maltose by KO11.

[0041] Figure 17 illustrates the results of the starch fermentation by recombinant *E. coli* strains KO11(pLOI140) and

KO11(pLOI141). (Since the results for both strains were essentially identical, only the data for KO11(pLOI141) is shown.)


[0042] Figure 18 comprises the thin layer chromatographic analysis of corn starch fermentation by KO11(pLOI141) during fermentation (72 hours).

[0043] Figure 19 comprises a schematic illustration of a fermentation process according to the invention.


Detailed Description of the Invention

I. Conventional Carbohydrate Metabolism

[0044] In most animals and plants as well as bacteria, yeasts and fungi, glucose is degraded initially by an anaerobic pathway prior to either oxidative or fermentative metabolism. The most common such pathway, termed glycolysis, refers to the series of enzymatic steps whereby the six-carbon glucose molecule is broken down, via multiple intermediates, into two molecules of the three carbon compound, pyruvate. During this process, two molecules of NAD⁺ are reduced to form NADH. The net reaction in this transformation of glucose into pyruvate is:




[0045] For glycolysis to continue, the NAD⁺ consumed by glycolysis must be regenerated by the oxidation of NADH. During oxidative metabolism, NADH typically is oxidized by donating hydrogen equivalents via a series of steps to oxygen and thereby forms water. Most organisms contain additional anaerobic pathways, however, which allow glycolysis to continue in the absence of compounds like oxygen. Such anaerobic processes are termed fermentation, and homolactic acid fermentation is perhaps one of the most common of these pathways occurring in many bacteria and in animals. In homolactic fermentation, glucose ultimately is converted to two molecules of the three carbon acid, lactic acid. NADH is oxidized by donating hydrogen equivalents to pyruvate and thus produces lactic acid. The net reaction for the regeneration of NAD⁺ by lactic acid fermentation is:



[0046] As mentioned previously, ethanologenic organisms like *Z. mobilis* and *S. cerevisiae* are capable of a second (alcoholic) type of fermentation whereby glucose is metabolized to two molecules of ethanol and two molecules of CO₂. Alcoholic fermentation differs from lactic acid fermentation in the steps used for the regeneration of NAD⁺. Two different enzymatic steps are required for alcoholic fermentation. Pyruvate decarboxylase cleaves pyruvate into acetaldehyde and carbon dioxide. Alcohol dehydrogenase serves to regenerate NAD⁺ by transferring hydrogen equivalents from NADH to acetaldehyde, thereby producing ethanol. The reactions for the regeneration of NAD⁺ by alcoholic fermentation are:



The net reaction for alcoholic fermentation is:



[0047] DNA that encodes each of these enzymes, respectively, had been cloned, and *Z. mobilis* pyruvate decarboxylase had been expressed recombinantly. See, for example, Brau & Sahm [1986a], *supra*. But it was highly uncertain whether the glycolytic pathway in organisms that do not produce ethanol as a primary metabolite ("non-ethanologenic organisms") could be diverted extensively at the pyruvate stage, even if such organisms could be modified somehow to express both of the additional enzymes required for alcoholic fermentation.

[0048] To the contrary, there was an expectation in the art that such a diversion might starve the recombinant host

for pyruvate and its normal products. See Brau & Sahm [1986a,b], *supra*. There also was a possibility that diversion of carbon from glycolysis to ethanol production in otherwise non-ethanogenic organisms would upset NAD/NADH ratios crucial to interrelated energy-generation and biosynthetic pathways.

5 II. Fermentation Pursuant to the Present Invention

[0049] It has been discovered, however, that organisms which carry out glycolysis or some variant thereof can be engineered, in accordance with the present invention, to divert as much as 95% of the carbon flow from glycolysis, at the pyruvate stage, to what in essence is a synthetic pathway comprised of the enzymes encoded by heterologous pyruvate decarboxylase (*pdc*) and alcohol dehydrogenase (*adh*) genes. The result is an engineered organism which produces ethanol as its primary fermentation product. That is, the ethanol produced exceeds 50% of the total fermentation product.

[0050] It also has been discovered that hosts can be selected for such heterologous expression which, by virtue of their native ability to transport and metabolize oligosaccharides, can ferment more complex feedstocks. (In this context, "oligosaccharide" denotes a molecule comprised of two or more saccharide monomers, including but not limited to the disaccharides cellobiose, maltobiose and xylobiose, trisaccharides like cellotriose and xylotriose, and long-chain polysaccharides such as cellulose, hemicellulose, starch, glycogen, pectin and inulin.) The capabilities of a host thus selected and transformed can be augmented, moreover, by expression in the same host of one or more genes coding for a polysaccharase, *i.e.*, an enzyme that catalyzes the breakdown to smaller oligosaccharides and/or saccharide monomers of complex oligosaccharides.

[0051] (A) **GENETICALLY ENGINEERING AN ARTIFICIAL ETHANOLOGENIC PATHWAY:** An important aspect of the present invention is an operon that directs cells to produce ethanol. Exemplary of such an operon is a construct of the present invention, designated a "*pet* operon," which comprises *Z. mobilis* genes coding for alcohol dehydrogenase II and pyruvate decarboxylase activities, respectively, together with appropriate regulatory sequences, for example, promoters, inducers, operators, ribosomal binding sites and transcriptional terminators. Thus, the dependence upon native alcohol dehydrogenase activity for ethanol production in previous recombinant systems is eliminated by combining *Z. mobilis* genes encoding alcohol dehydrogenase II and pyruvate decarboxylase to form a *pet* operon. Moreover, significant amounts of ethanol can be produced, in recombinants containing the *pet* operon, under both aerobic and anaerobic conditions.

[0052] To impart to a microorganism the ability to produce polysaccharases, such as xylanases and cellulases, a ethanogenic operon of the present invention can be modified by adding the gene(s) which code for the desired enzyme(s). Alternatively, one or more polysaccharase-encoding genes can be incorporated into a plasmid which is used to transform a host organism already engineered with an operon that directs ethanol production, as described above. (This aspect of the present invention is described in more detail in Section II(E) below.) By yet another alternative approach, the host to be transformed with the ethanogenic operon is selected for its native ability to express a polysaccharase. Yet another approach is to add the polysaccharase genes into the chromosome by integration.

[0053] The approach of combining, in a single construct, the genes that encode a metabolic pathway is generalized readily to a variety of situations where genes from different loci can be brought together to create an artificial operon. The creation of a *pet* operon thus is but one example of the applications envisaged for the present invention. For example, genes coding for alcohol dehydrogenase activity from a variety of organisms could be combined with other genes coding for pyruvate decarboxylase activity in order to create an operon coding for the desired pathway. Operons coding for other pathways also could be created. It also is apparent that, for the above-discussed ethanogenic pathway, it is not necessary that the genes coding for alcohol dehydrogenase and pyruvate decarboxylase activities be under common control; they could be under separate control, even in different plasmids or in different locations on the chromosome. Likewise, gene(s) encoding polysaccharases could be under common or separate control, or located in different plasmids or at different chromosomal positions.

[0054] Another aspect of the present invention concerns the use of recombinant ethanol-producing hosts for the efficient production of recombinant peptides or proteins; that is, the recombinant cells can be transformed further with genes coding for useful products other than polysaccharases. (These additional genes could be plasmid-borne or incorporated into the chromosome.) More specifically, genes that encode the necessary ethanogenic enzymes generally are expressed at high levels and dominate carbon flow from pyruvate and NADH oxidation during anaerobic growth. Under these conditions, the flow of pyruvate carbon skeletons can be diverted from the production of organic acids to the production of ethanol as the principal fermentation product.

[0055] In this way, the extent of acidification per unit of cell protein is minimized by the production of ethanol rather than of organic acids. Oxygen transfer is frequently a major limitation during the growth of dense cultures of microorganisms, and it is this limitation which results in acid production and pH drift of the growth medium. In recombinants expressing an ethanogenic operon within the present invention, by contrast, the heterologous, ethanogenic enzymes divert part of the pyruvate from glycolysis to acetaldehyde and reoxidize NADH to produce ethanol, a less

damaging product of metabolism. Thus, strains containing both functional respiratory chains for oxidative phosphorylation and ethanol production enzymes can be grown to even higher cell densities because of the operation of both systems during the regeneration of NAD⁺ and a reduction in acidic waste products. Such inherent flexibility results in less stringent process-control requirements, as well as increased volumetric yields of recombinant products.

[0056] The accumulation of organic acids which otherwise occurs is regarded as a consequence of fermentation during anaerobic growth. But appreciable quantities of acetate may be produced even under aerobic conditions, e.g., during rapid agitation. Thus, the production of acetate may be progressive from the earliest stages of growth and not limited to the later stages, when cell density is high and anaerobic conditions prevail. Acid production from glucose, even under aerobic conditions, serves to limit growth in broth and on solid medium, as demonstrated by the increased final cell density in medium supplemented with phosphate buffer.

[0057] Accordingly, ethanol-producing transformants within the present invention also are superior hosts for production of recombinant products, even under anaerobic conditions, with minimal acid production. Many recombinant proteins and peptides contain cysteine or disulfide bridges, and proper folding or reactions of these is an essential feature to form the active enzyme. Since formation of disulfide bonds is promoted by oxygen, synthesis of such proteins under anaerobic conditions provides less opportunity for improper folding prior to isolation and folding under controlled conditions, thus resulting in greater recovery of biologically active product.

[0058] The use of *adhB* in these constructs may be of particular advantage. *AdhB* encodes a stress protein (An *et al.* [1991] *FEBS LETTERS* 282:205-208). Stress proteins have been shown to aid in the proper folding of heterologous proteins allowing the retention of biological function. Lee *et al.* [1992] *J. Biol. Chem.* 267:2849-2852. Accordingly the use of *adhB* could enhance the production of functional proteins in recombinant organisms.

[0059] Under aerobic conditions in *E. coli*, pyruvate from glycolysis is primarily metabolized by the pyruvate dehydrogenase complex and by lactate dehydrogenase, with excess acetyl coenzyme A being converted to acetate. See Gottschalk, *BACTERIAL METABOLISM*, pp. 210-80 (Springer-Verlag 1986). The apparent K_m values for these two enzymes are 0.4 and 7.2 mM, respectively. The apparent K_m for *Z. mobilis* pyruvate decarboxylase is equal to (pyruvate dehydrogenase) or lower than (lactate dehydrogenase) those for the two *E. coli* enzymes, thereby facilitating acetaldehyde production. NAD⁺ regeneration under aerobic conditions results primarily from biosynthesis and from the NADH oxidase (coupled to the electron transport system) with an apparent K_m of 50 μM. The apparent K_m for *Z. mobilis* alcohol dehydrogenase II is over four-fold lower than that for *E. coli* NAD⁺ oxidase, allowing the *Z. mobilis* enzyme to compete effectively for endogenous pools of NADH for the reduction of acetaldehyde to ethanol. Accordingly, the properties of the *Z. mobilis* ethanologenic enzymes and their relatively high levels of expression are well suited for the diversion of carbon flow into ethanol under aerobic conditions.

[0060] Under anaerobic conditions in *E. coli*, pyruvate from glycolysis is primarily metabolized by lactate dehydrogenase and pyruvate formate lyase. The apparent K_m values for these two enzymes are 18-fold and 5-fold higher, respectively, than that for *Z. mobilis* pyruvate decarboxylase. Similarly, the apparent K_m's for the principal enzymes involved in NAD⁺ regeneration in *E. coli* also are considerably higher than those for *Z. mobilis* alcohol dehydrogenase II. Thus, the ethanologenic enzymes from *Z. mobilis* are quite competitive for carbon (pyruvate) and reducing potential (NADH) with the normal fermentative enzymes and oxidative phosphorylation in *E. coli*, allowing the efficient channeling of glycolytic products into ethanol.

[0061] It has been discovered that lactose and all of the major sugars present in cellulose and hemicellulose (namely, glucose, xylose, arabinose, galactose and mannose) can be converted to ethanol, in accordance with the present invention, by recombinant hosts that express genes encoding ethanologenic enzymes, such as those comprising the above-discussed ethanol pathway of *Z. mobilis*.

[0062] A variety of factors should be considered in selecting host strains suitable for ethanol production pursuant to the present invention. These factors include substrate range and environmental hardiness, such as sugar tolerance, salt tolerance, ethanol tolerance, tolerance to low pH, and thermal tolerance. As described below, for example, *E. coli* strain ATCC 9637 (Waksman strain W) displays superior characteristics in terms of environmental hardiness, although ethanol production from glucose was lower than with other strains. Strain ATCC 9637 was selected primarily for its unique ability to utilize sucrose. Advantageously, this characteristic of ATCC 9637 makes this strain useful in the fermentation of beet sugar, cane sugar, and other feedstocks with sucrose. ATCC 11303 (Luria strain B) and ATCC 15244 (Kepes strain ML300) containing pLOI297 produced the highest levels of ethanol and exhibited acceptable levels of environmental hardiness. Plasmids were quite stable in these two constructs and they were selected as the best candidates for further development of ethanol production. Both constructs expressed the high levels of *Z. mobilis* pyruvate decarboxylase which are required for efficient ethanol production. See Ingram *et al.* [1987].

[0063] All major sugar components of plant biomass were converted to ethanol by the recombinant *E. coli* containing the ethanol pathway from *Z. mobilis*. The conversion efficiency of glucose and xylose into ethanol exceeded that which has been previously reported for *S. cerevisiae*, see Lovitt *et al.* [1988] *CRC Crit. Rev. Biotechnol.* 7: 107-186, and for pentose-fermenting yeasts systems. See Beck [1989] *Biotechnol. Bioeng. Symp.* 17: 617-627; Jeffries *et al.* [1988] *Biotechnol. Bioeng.* 31: 502-506; Skoog *et al.* [1988] *Enzyme Microbiol. Technol.* 10:66-79; Slininger *et al.* [1985]

Biotechnol. Lett. 7:431-436). Xylose was converted to ethanol by recombinant *E. coli* with a higher efficiency than glucose by *S. cerevisiae*, as reported by Lovitt *et al.*, *supra*. The unusually high ethanol yields with xylose (over 100% of theoretical) might be attributed to ethanol derived from the catabolism of complex nutrients. Many amino acids and complex medium components are catabolized to glycolytic intermediates or to pyruvate directly. This pyruvate could then be converted to ethanol.

[0064] According to the present invention, *pdh* and *adh* genes can be transformed into a variety of different hosts and expressed using a variety of promoters. It is well within the skill of a person trained in this field to use the descriptions provided herein to make these transformations. For example, *pdh* and *adh* genes can be readily inserted into plasmids which have different host ranges. Plasmids which are constructed to replicate in *E. coli* and another target organism are typically called "shuttle vectors," since they are able to replicate in two or more hosts. These plasmids are readily available and easily used. Most frequent combinations include shuttles between *E. coli* and other bacteria, *E. coli* and yeasts, and *E. coli* and animal cells. These vectors are available from catalogs and are well known to those skilled in the art.

[0065] *Z. mobilis* *pdh* and *adh* genes have been inserted and expressed in *Xanthomonas*, *Klebsiella* and *Erwinia*. Similar success in other organisms also can be achieved readily. Although some straightforward modification of the promoter, for example, may be needed to produce the optimal construct, success is highly predictable based upon biochemical properties and current knowledge of expression of soluble cytoplasmic proteins. Foreign genes expressing enzymatic activity needed to redirect pyruvate metabolism, in accordance with the present invention, can even be integrated into the chromosome and expressed without the need for a plasmid. See Section II(C), *infra*. For example, a pet construct which lacks a promoter can be integrated into the *E. coli* chromosome immediately behind the promoter for the pyruvate formate lyase gene. Analogous integration into *pfl* or other genes is possible in most organisms, requiring only a fragment of the target gene to direct the site of integration by homologous recombination. Target genes other than *pfl* also could be used for integration. Since pet-expressing constructs are easily identified on indicator plates, this approach is widely feasible for the rapid and efficient construction of other organisms for ethanol production.

[0066] A great many texts are available which describe procedures for expressing foreign genes. Also, catalogs list vectors which can be used for various organisms. Cloning vectors for gram-negative bacteria, gram-positive bacteria, vectors with broad host range capability, fungal cloning vectors, insect cloning vectors, animal cell cloning vectors, and plant cell cloning vectors are all well known to those skilled in the art. Catalogs from which these cloning vectors can be ordered and are readily available and well known to those skilled in the art. See, for example, Marino [1989] *BioPharm* 2: 18-33; VECTORS: A SURVEY OF MOLECULAR CLONING VECTORS AND THEIR USES (Butterworths 1988).

[0067] The skilled practitioner also has access to alternative *pdh* and *adh* genes, and to other such genes which can be identified by use of the aforementioned *Z. mobilis* genes as probes or, more preferably, by observing activity on indicator plates. For purposes of this invention, it does not matter whether the alcohol dehydrogenase activity is provided from a gene isolated from a horse, yeast, human, insect, or other bacterial gene. Since expression of alcohol dehydrogenase activity can be observed directly on aldehyde indicator plates, sequence information would not be needed for the isolation of additional genes encoding proteins which exhibit this enzymatic activity. Indeed, many alcohol dehydrogenase genes are already well known to those skilled in the art, as evidenced by the recitation of 252 *adh* genes in the Genbank database as of March 1991 (IntelliGenetics Inc., 700 E. El Camino Drive, Mountain View, CA 94040).

[0068] *Z. mobilis* contains two genes encoding functional alcohol dehydrogenase genes, and one of these (*adhB*) is related evolutionarily to a butanol dehydrogenase from *Clostridium acetobutylicum*, propanediol oxidoreductase from *E. coli*, and ADHIV alcohol dehydrogenase from *Saccharomyces*. All have been cloned and sequenced. The second alcohol dehydrogenase gene from *Z. mobilis*, *adhA*, is a zinc alcohol dehydrogenase which also has been cloned and sequenced. Based upon comparisons of primary structure deduced from sequences which are readily available, the *adhA* gene is deemed related evolutionarily to the typical alcohol dehydrogenases described in animals, plants, and the dominant *adh* gene in yeasts. The *adhA* gene and other alcohol dehydrogenase genes can be substituted for the original *adhB* gene exemplified herein.

[0069] By the same token, for purposes of this invention it does not matter whether the pyruvate decarboxylase activity is provided by a gene from *Z. mobilis* or by a gene which encodes the needed enzymatic activity but which comes from corn, yeast or some other organism. The evolution of life forms from common ancestry is now well accepted and has been demonstrated in splendid detail by the methods of molecular genetics. Not only can organisms be arranged in phylogenetic trees based upon macro-characteristics, but the ancestral genes which have evolved for specific functions have been retained throughout evolution with conservation of features required for such functions. This high level of conservation enables those skilled in the art to isolate functionally equivalent, genetically related enzymes from other organisms using primary information from one or more members of an enzyme family. The enzymes of glycolysis are some of the best examples of this since such enzymes have been so well studied.

[0070] Indeed, just such an approach has been used successfully to clone the pyruvate decarboxylase gene from maize, using information on the *Z. mobilis* *pdh* and the *pdh* of *S. cerevisiae* to design a DNA probe. See Kelly [1989] *Plant Molecular Biology* 13: 213-22. Alternative strategies could have used the entire genes as probes. Since the

synthesis of a protein with pyruvate decarboxylase activity (pyruvate converted to acetaldehyde plus carbon dioxide) can be observed directly on aldehyde indicator plates, sequence information would not be needed to locate other genes, although sequence information has been used to isolate the corn gene. Many other pyruvate decarboxylase genes which provide a functional equivalent can be isolated from other organisms. These other genes would be suitable replacements for the *Z. mobilis pdc*, just as several alcohol dehydrogenases have proven suitable. Further in that regard, the GenBank database listed at least 5 *pdc* genes as of March 1991.

[0071] As a further improvement, a recombinant host within the present invention can be transformed with genes encoding key components of the glucose transport in *Z. mobilis*, namely, the genes coding for *Z. mobilis* glucokinase (*glk*) and for glucose-facilitated diffusion protein (*glf*). In particular, *glk* and *glf* can be combined in an artificial operon, along the lines described above for an ethanologenic operon such as *pet*, and transformed into a recombinant already containing the latter operon. Expression of the *glk/glf* operon would result in an increase in trans-membrane glucose flux and, hence, in an increased rate of ethanol production. Likewise, genes encoding key components of the transport of oligosaccharides in other hosts, or genes encoding key components in the glycolytic pathway, also can be used to further transform hosts in accordance with this invention. Such transformation with genes encoding rate-limiting steps in flux can be expected to increase the rate of ethanol production and boost the levels of glycolytic intermediates as a source of skeletons for biosynthesis.

[0072] (B) **FERMENTATION PARAMETERS:** Pursuant to the present invention, heterologous ethanologenic genes are expressed at levels that facilitate the production of ethanol as a fermentation product by the recombinant host. By inserting such genes into an appropriate non-ethanologenic organism, as described above, the transformed host is enabled to produce significant amounts of ethanol from oligosaccharides like cellobiose, cellotriose, xylobiose and xylotriose. As discussed below, ethanol concentrations on the order of 1 M, and up to 1.5 M can be obtained from recombinant hosts according to the present invention.

[0073] (C) **CHROMOSOMAL INTEGRATION OF FOREIGN GENES:** Chromosomal integration of foreign genes can offer several advantages over plasmid-based constructions, the latter having many limitations for commercial processes. Plasmid-based constructions are inherently less stable than chromosomal genes and serve as an environmental hazard, a reservoir for transmission of heterologous traits. Instability often is exacerbated by the addition of multiple plasmids. Nevertheless, plasmid-based constructions can be useful for initial experiments involving expression of heterologous genes, such as genes encoding cellulases, since they can be used to readily transform host cells. But to facilitate screening of the resulting protein (e.g., a cellulase), it is often desirable to eliminate the need for a plasmid carrying ethanol-producing genes by chromosomal integration in accordance with this invention.

[0074] Ethanologenic genes have been integrated chromosomally in *E. coli* B, see Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57: 893-900. In general, this is accomplished by purification of a DNA fragment containing (1) the desired genes upstream from a chloramphenicol gene and (2) a fragment of homologous DNA from the target organism. This DNA can be ligated to form circles without replicons and used for transformation. Thus, the *pfl* gene can be targeted in the case of *E. coli*, and short, random *Sau3A* fragments can be ligated in *Klebsiella* to promote homologous recombination.

[0075] Initial selections of recombinants can be made on 20 mg chloramphenicol ("Cm")/liter plates to allow growth after single copy integration. These constructs may be obtained at a very low frequency. Ethanologenic genes initially may be expressed at low levels, insufficient to permit efficient ethanol fermentation. Higher level expression may be achieved as a single step by selection on plates containing 600 to 1000 mg Cm/liter. Such strains have proven very stable and it is anticipated that the same approach will be successful for more optimal strains of *Klebsiella* and *Erwinia*. Preliminary testing of a few wild strains indicates that electroporation improves plasmid delivery and may reduce the effort required to achieve integrations. (See Example 16 below.)

[0076] (D) **HOST SELECTION:** The range of organisms suitable for modification to express heterologous *pdc* and *adh* genes, as described above, includes, *inter alia*, eukaryotic cells, such as animal cells, insect cells, fungal cells, and yeasts which are not naturally ethanologenic, and non-ethanologenic bacteria. In addition to *E. coli*, for example, other enteric bacteria of the genera *Erwinia*, like *E. chrysanthemi*, and *Klebsiella*, like *K. planticola* and *K. oxytoca*, are particularly attractive hosts because they can metabolize a variety of sugars, including pentoses. Advantageous hosts can also be selected from the broader category of gram-negative bacteria, such as species of the genus *Xanthomonas*, and from the gram-positive bacteria, such as members of the genera *Bacillus*, *Clostridium* and *Cellulomonas*. Appropriate transformation methodology is available for each of these different types of hosts.

[0077] Certain organisms among the aforementioned microbes also meet the criteria for selection of a host to ferment oligosaccharide(s) to ethanol in accordance with the present invention. More specifically, a host can be selected in this regard because it produces (1) the proteins necessary to transport an oligosaccharide into the cell and (2) intracellular (cytoplasmic) levels of enzymes which metabolize those oligosaccharides. Microbes mentioned previously that meet these criteria include enteric bacteria like *E. chrysanthemi* and other *Erwinia*, and *Klebsiella* species such as *K. oxytoca*, which naturally produces a β -xylosidase, and *K. planticola*. Certain *E. coli* also satisfy the criteria because they transport and metabolize cellobiose, maltose and/or maltotriose. See, for example, Hall *et al.* [1987] *J. Bacteriol.* 169: 2713-17.

[0078] Hosts can be selected, in satisfaction of criteria (1) and (2) above, from the broader categories of gram-negative bacteria, such as *Xanthomonas* species, and gram-positive bacteria, including members of the genera *Bacillus*, such as *B. pumilus*, *B. subtilis* and *B. coagulans*; *Clostridium*, for example, *Cl. acetobutylicum*, *Cl. aerotolerans*, *Cl. thermocellum*, *Cl. thermohydrosulfuricum* and *Cl. thermosaccharolyticum*; *Cellulomonas* species like *C. uda*; and *butyrivibrio fibrisolvens*. Similarly acceptable are various yeasts, exemplified by species of *Cryptococcus* like *Cr. albidus*, species of *Monilia*, *Pichia stipitis* and *Pullularia pullulans*; and other oligosaccharide-metabolizing bacteria, including but not limited to *Bacteroides succinogenes*, *Thermoanaerobacter* species like *T. ethanolicus*, *Thermoanaerobium* species such as *T. brockii*, *Thermobacteroides* species like *T. acetohydrolyticus*, and species of the genera *Ruminococcus* (for example, *R. flavefaciens*), *Thermonospora* (such as *T. fusca*) and *Acetivibrio* (for example, *A. cellulolyticus*). Again, suitable transformation techniques are available for these different hosts.

[0079] The literature relating to microbes which meet the subject criteria is reflected, for example, in Biely [1985] *Trends in Biotech.* 3: 286-90, in Robsen *et al.* [1989] *Enzyme Microb. Technol.* 11: 626-44, and in Beguin [1990] *Ann. Rev. Microbiol.* 44: 219-48, the respective contents of which are hereby incorporated by reference. Those skilled in this art will appreciate that many other hosts will be suitable for use in the present invention. Thus, suitable hosts can be identified by screening to determine whether the tested microbe transports and metabolizes oligosaccharides. Screening in this vein could be accomplished in various ways. For example, microorganisms could be screened to determine which grow on suitable oligosaccharide substrates, the screen being designed to select for those microorganisms that do not transport only monomers into the cell. See, for example, Hall *et al.* [1987], *supra*. Alternatively, microorganisms could be assayed for appropriate intracellular enzyme activity, *e.g.*, β -xylosidase activity.

[0080] Exemplary of a selection regimen suitable for use in accordance with the present invention is one where minimal medium containing pectin is employed to isolate a soft-rot *Erwinia* strain from decaying plant materials. See Starr, "The Genus *Erwinia*," in Vol. II THE PROKARYOTES 1260-71 (Springer-Verlag 1981). Isolates which cause pitting in calcium stabilized pectin can be characterized further to confirm identity. See Bergey's Manual, Volume 1, at pages 469-76.

[0081] Suitable candidate strains of *Klebsiella*, abundant in pulp and paper waste (Huntley *et al.* [1976] *J. Water Pollution Control Federation* 48: 1766-1771; Knittel *et al.* [1977] *Appl. Environ. Microbiol.* 34: 557-63), can be obtained, for example, from waste streams of processing plants in Perry and Palatka, Florida. The presence of high numbers of *Klebsiella* in such waste may promote selection for particularly useful strains with optimal characteristics such as resistance to toxic chemicals and the ability to utilize oligosaccharides. Strains can be selected initially by growth on coliform-specific medium, see Seidler, "The Genus *Klebsiella* (Nonmedical Aspects)," in II THE PROKARYOTES 1166-72, and further identified by indole-production, Voges-Proskauer reaction, citrate, etc. See Bergey's Manual, Volume 1, at pages 461-65.

[0082] Isolates of *Klebsiella* and *Erwinia* can be compared for their ability to degrade carboxymethyl cellulose, xylan-blue, methyl-umbelliferyl β -D-glucopyranoside, methyl-umbelliferyl cellobioside, methyl-umbelliferyl xyloside, and methyl-umbelliferyl arabinoside as model substrates for endoglucanase, xylanase, β -glucosidase, cellobiohydrolase, xylosidase and arabinosidase, respectively, on plates buffered at pH 6.0 (30°C). Strains also should be screened for their ability to metabolize monosaccharides (hexoses and pentoses) as well as cellobiose, cellobiose, xylobiose and xylobiose. Disaccharides and trisaccharide levels can be tested in overnight tube cultures by means of thin layer chromatography to monitor utilization of saccharides during overnight incubation. Xylosidases can be generated from birchwood xylan (Sigma) by treatment with *C. thermocellum* xylanase.

[0083] Growth of promising strains can be further screened for ethanol tolerance, salt tolerance, and temperature tolerance in a manner essentially similar to that used for the selection of *E. coli* B. See Alterhum *et al.*, [1989] *Appl. Environ. Microbiol.* 55: 1943-48; Beall *et al.*, *Biotechnol. & Bioeng.* [1991] 38: 296-303. Although most isolates can be frozen in liquid nitrogen for storage, the three most promising strains from each group can be selected for further development. Stored strains may prove useful in future studies as sources of new polysaccharase genes.

[0084] Optimal strains should retain the ability to efficiently metabolize all constituents of lignocellulose (disaccharides, trisaccharides and monosaccharides), to grow in ethanol concentrations above 440 g/liter, to tolerate salt levels of 0.3 molar, to tolerate acetate levels of 0.2 molar, and to tolerate temperatures of 40°C and to produce high levels of enzymes useful for cellulose, hemicellulose and pectin depolymerization with minimal protease activity. Some strains may also contain native xylanases or cellulases. After integration of foreign genes for alcohol production, optimal constructions should produce ethanol from all saccharides tested with greater than 90% of theoretical yield while retaining the useful traits above.

[0085] Additional studies can be done to evaluate overall pH and temperature optima of secreted cellulases and to determine thermal stability. This information can be helpful in designing optimal conditions for combined hydrolysis and fermentation processes.

[0086] (E) EXPRESSION OF HETEROLOGOUS POLYSACCHARASES: As described above, a preferred embodiment of the present invention contemplates production of recombinants which express heterologous polysaccharases, thereby reducing the need for commercial enzymes in the context of fermenting oligosaccharides. Exemplary polysac-

charases include cellulolytic enzymes, hemicellulytic enzymes, xylanolytic enzymes and starch-degrading enzymes.

[0087] In one advantageous embodiment, the polysaccharase is thermostable. In this regard, for example, genes encoding thermophilic proteins from *Clostridium thermocellum* (*celA*, *B*, *C* and *D*) can be expressed individually at high levels as cytoplasmic products. Recombinantly reproduced enzymes are recovered by harvesting cells at the end of fermentation and then releasing the thermostable enzymes from those cells by heating, for example, to 60°C, a temperature at which they exhibit near maximal activity and good stability. Thereafter, the enzymes are allowed to act on a complex polysaccharide substrate for a predetermined period, e.g., on the order of twelve hours. Modifications which eliminate the signal portions of selected genes greatly enhance the activities which are produced in *E. coli*, and may have a similar effect in *Klebsiella* and *Erwinia*.

[0088] The *celD* gene is particularly preferred for use in this context, and *celD*-transformed recombinants within the present invention are suitable for the fermentation of a substrate like SOLKA-FLOC, a powdered cellulose product derived by mechanical comminution of purified wood pulp. As shown in Example 18 and Figure 14 below, prehydrolysis of substrate with the *celD*-encoded enzyme can reduce the requirement for commercial cellulases to approximately 1/5 in relation to recombinants which lack *celD*. This approach of using enzymes, heterologously expressed and accumulated intracellularly, to "jump-start" saccharification of a complex substrate can be generalized, in accordance with the present invention, to any ethanologenic recombinant that is further transformed with a polysaccharase-encoding gene, the expression product of which can be recovered for the purpose of breaking down complex oligosaccharides (cellulose, hemicellulose, starch, glycogen, pectin, inulin, etc.). See Examples 17-19 below.

[0089] Thus, for example, cells from a previous fermentation can be used as a source of enzymes for each subsequent fermentation. Also, enzyme secretion is not required and, instead, the process takes advantage of intracellular accumulation to facilitate harvesting of enzymes. The highest level of recombinant enzyme is present initially rather than accumulating slowly during growth and fermentation. Furthermore, there is reduced risk of microbial contamination since oligosaccharide utilization is uncommon among microorganisms and, hence, fewer contaminating microflora are to compete for these sugars.

[0090] In another preferred embodiment, an ethanologenic recombinant as described above expresses a polysaccharase that is at least partially secreted from the host and catalyzes hydrolysis of a complex oligosaccharide extracellularly. Such a gene product could supplement the activity of one or more commercial enzymes, such as the products of *cenA* (endoglucanase) and *cex* (exoglucanase), conventionally supplied during saccharification and ethanol production. Illustrative of polysaccharase-encoding genes which could be used in this matter are the cellulase genes of *Cellulomonas fimi*, which are partially secreted (about 50% to 60%) when transformed into *Klebsiella* or *Erwinia*. In a laboratory strain of *Erwinia* transformed with the *cex* gene, for example, over 50% of the heterologous expression product is secreted into the broth.

[0091] The efficiency of such secretion should be increased by fusion to signal sequences for pullulanase in *Klebsiella* and for a pectate lyase in *Erwinia*, respectively. See Murooka [1990] *Ferment. Technol. Ind. Appl.* 40-45; Pugsley et al. [1990] *Ann. Rev. Genetics* 24: 67-90; He et al. [1991] *Proc. Nat'l Acad. Sci. USA* 88: 1079-83; Saarilahti et al. [1990] *Gene* 90: 9-14.

[0092] Of particular interest is an ethanologenic transformant within the present invention that expresses both a thermostable gene product in addition to a gene product which is secreted. Such a transformant would provide (1) via lysis, enzymatic activity for use in a pre-treatment to break cellulose or another complex substrate into simpler oligosaccharides and monosaccharides, and (2) via secretion, enzymes for continued depolymerization during fermentation.

[0093] Heterologous genes coupled with mutations or genetic modifications that overexpress native traits will prove useful under particular circumstances. For example, mutageneses and selection can be carried out to increase the level of expression of native endoglucanases in *Erwinia* strains. In some cases these are repressed by high concentrations of glucose. Selection for depressed mutants on glucose-CMC plates can be used to identify strains which have lost this control, some of which may over-produce these enzymes.

[0094] The skilled artisan will appreciate that many different polysaccharases can be employed in accordance with this aspect of the present invention. By the same token, a wide range of transforming vectors are available for use in accordance with this aspect of the present invention. Some routine experimentation may be necessary to achieve the optimal match of host and vector. For example, RSF1010-based vectors can be constructed, see Conway et al. [1987c] *Appl. Environ. Microbiol.* 53: 235-41, and are preferred to the extent that their broad host range should allow introduction of heterologous genes into many gram-negative bacterial host, including all *Erwinia* and *Klebsiella* isolates, with selection for tetracycline resistance. By contrast, pUC-based plasmids may prove unstable in some *Erwinia* strains and, to that extent, would be less preferred.

[0095] Among other polysaccharase genes suitably used in the present invention are those that encode one of the cellulolytic activities listed below:

Type of Enzyme	Substrate				
	Crystalline cellulose	Carboxymethyl Cellulose (CMC)	Amorphous Cellulose	Cellobiose	Cellobiose
Endoglucanase	-	+	+	+	-
Cellobiohydrolase	+	+/-	+	+	-
β -Glucosidase	-	-	-	+	+

[0096] An endoglucanase hydrolyses β -1,4-glycosidic linkages randomly. It does not attack cellobiose but hydrolyzes cellobioextrins, phosphoric-acid-swollen cellulose and substituted celluloses like CMC and HEC (hydroxyethyl cellulose). Some endoglucanases also act on crystalline cellulose.

[0097] By contrast, a cellobiohydrolase acts on cellulose and, in particular, splits off cellobiose units from the non-reducing end of the chain. Cellobiohydrolase hydrolyses cellobioextrins but not cellobiose. β -Glucosidase hydrolyses cellobiose and cellobiooligosaccharides to glucose, but does not attack cellulose or higher cellobioextrins.

[0098] In addition to the cellulase-encoding genes from *C. thermocellum* mentioned above, other genes that code for cellulolytic activity have been characterized. See, for example, Bèguin [1990] *Ann. Rev. Microbiol.* 44: 219-48, the contents of which are hereby incorporated by reference. Moreover, such genes are readily available, for example, through GenBank database, which contained listings of 13 endoglucanases, 3 cellobiohydrolases, and 3 β -glucosidases, *inter alia*, as of March 1991.

[0099] By the same token, genes that code for a xylanolytic enzyme can be used as described above, pursuant to the present invention. Of particular interest in this regard are genes encoding a xylanolytic enzyme selected from the group of endo-1,4- β -xylanase, β -xylosidase, α -glucuronidase, α -L-arabinofuranosidase, acetylsterase and acetylxy-lanesterase. Exemplary of this group are the respective expression products of the *C. thermocellum xynZ* gene and the *Butyrivibrio fibrisolvens xylB* gene, both expressed in ethanologenic transformants described below (see Example 17). Again, genes for such xylanases are readily available as reflected in GenBank's entries for 7 endo-1,4- β -xylanases, 5 β -xylosidases and 1 α -L-arabinofuranosidase as of March 1991.

[0100] Similarly, genes can be used in the present invention that code for enzymes involved in the breakdown of starch. Illustrative enzymes of this sort are listed below, along with biological source and other characterizing information:

Enzyme	Type	Source	Amount	Activity
α -Amylase	Liquefying	<i>Bacillus subtilis</i> ^a	0.06% wt of starch	Decreases viscosity (Cleaves 1-4, pH 5.5, 70°C)
		<i>B. licheniformis</i>	0.06% wt of starch	Decrease viscosity (92°C)
		Barley malt	0.5-1.0 wt of grain	Decreases viscosity (60°C)
β -Amylase	Saccharifying	Barley malt ^b	2.0% wt of grain	Generates maltose (Cleaves 1-6, pH 5.5, 60°C)
Glucoamylase	Saccharifying	<i>Aspergillus niger</i> ^b	0.18% wt of starch (1.7 L/ton)	Generates glucose (Cleaves 1-6, pH 5.0, 60°C)
Pullulanase	Saccharifying	<i>B. acidopullulyticus</i>	0.2% wt of starch (2.0 kg/ton)	Cleaves 1-6, pH 5.0, 60°C

^a Endoamylase

^b Exoamylase

[0101] Enzymes involved in starch breakdown are exemplified below by a *B. stearothermophilus* α -amylase and a *T. Brockii* pullulanase. Such genes are likewise readily available, as reflected by GenBank's entries, as of March 1991, for 190 α -amylases, 3 β -amylases, 8 glucoamylases, and 3 pullulanases. Other polysaccharases such as β -glucanase,

arabinosidase, mannanase, pectin hydrolase, and pectate lyase also are available readily through GenBank or from literature which teaches the cloning of such genes.

[0102] *Bacillus* polysaccharase genes often are expressed well in *E. coli* and, hence, should be suitable for use in enteric bacteria generally, pursuant to the present invention. Indeed, members of the Bacillaceae secrete a wide variety of such enzymes, including amylases, β -glucanases and hemicellulases, and a number of species secrete cellulases, including strains of *B. subtilis*, *B. polymyxa*, *B. licheniformis* and *B. cereus*. See Bèguin [1990], *supra*, and Robson *et al.* [1989] *Enzyme Microb. Technol.* 11: 626-44, the contents of which are hereby incorporated by reference. If a signal sequence associated with a heterologous polysaccharase gene proves non-functional in the host of choice, such as *Klebsiella* or *Erwinia*, it is a routine expedient to replace this signal by gene fusion to a signal sequence from the host in the event that, as described above, secretion of heterologous enzyme is desired.

[0103] Strains of *Bacillus stearothermophilus* are a source of genes encoding thermostable hydrolase and other activities, such as endoglucanase (CMCase and nitrophenyl-cellobiosidase activities), which can be exploited to advantage in the present invention. Many genes of this type have been characterized, and are available through GenBank or from the relevant literature. See also Bèguin [1990], *supra*.

[0104] (F) SELECTION AND TRANSFORMATION OF HOSTS: To select strains as potential hosts for heterologous polysaccharase genes, recombinant strains within the present invention which contain integrated genes for alcohol production can be tested in fermentations using SOLKA-FLOC as the cellulosic substrate, together with a commercial cellulase for hydrolyzing the cellulose. A dose-response curve can be generated for ethanol production and rate of conversion as functions of dilution of commercial enzyme preparation. The industrial cellulase preparations (e.g., SPEZYME, MULTIFECT AND CYTOLASE [Genencor]) can be used in these experiments, for example, using pH-controlled fermentations at 35°C and pH 6.0.

[0105] Addition of a heterologous polysaccharase then can be accomplished in the manner illustrated in Examples 17-19. For example, polysaccharase genes which are selected as particularly advantageous can be integrated using homologous recombination. Tetracycline (5 mg/liter) can be used for selection of initial recombinants and higher levels will be used to select mutants with elevated expression of useful genes. Fusaric acid selection can be used to inactivate the tetracycline resistance (*tet*) gene so that multiple enzymes can be added sequentially by the same protocol. These constructs can be evaluated for utility in the single stage and multistage cellulose fermentation processes.

[0106] A preferred route to developing ethanologenic, gram-positive organisms, such as a *B. subtilis* or a *Cellulomonas* strain, involves chemical mutagenesis coupled with screening of colonies by the use of antibodies against *Z. mobilis* pyruvate decarboxylase, thereby to detect increases in expression. Clones thus identified for displaying higher expression are studied to determine the mechanism(s) responsible for the phenotype. Such mutants may prove useful *per se* and, in addition, should serve well for the expression of other heterologous proteins in an organism which is generally regarded as safe (GRAS).

[0107] In transforming ethanologenic hosts with heterologous cellulase genes, thereby to achieve high intracellular concentrations of active enzyme, some additional, routine genetic manipulation or screening may be required, depending upon the host and cellulase selected. For example, proteases secreted by a host may present a barrier to heterologous expression in accordance with the present invention. Several approaches can be used, such as indicator plates for proteinase activities and selection for higher levels of target enzyme, to obtain mutants in which this problem has been minimized.

[0108] Moreover, there often is a price to pay, in terms of growth rate and efficiency, for the production of any heterologous protein, including a polysaccharase. This "gene burden" can be minimized by testing multiple constructions of particularly useful genes. The site of integration also may affect the apparent "gene burden" and, therefore, can be a variable for testing in accordance with the present invention.

[0109] The constructs prepared along these lines then can be used in cellulose fermentation trials, as illustrated in Example 18 below, to evaluate effectiveness in combination with commercial enzymes and for efficiency of fermentation. Thus, one can test the ability of cellulases that are secreted by the transformed host to replace commercial cellulases in single-stage processes where commercial cellulases are added at the time of inoculation with ethanol-producing, cellulase-secreting recombinants. The utility of thermophilic cellulases can be tested by adding ethanologenic cells containing these enzymes from a prior fermentation to cellulose suspensions at 60°C. Incubation can be continued for a period of time, such as 12 hours, under optimized conditions for the particular enzyme. The temperature and pH can be lowered in most cases to provide optimal conditions for a commercial enzyme, such as SPEZYME, MULTIFECT and CYTOLASE, and incubation continued for 12 hours. After lowering the temperature to 35°C, and raising the pH to 6.0, broth can be inoculated with recombinant ethanologen. Ethanol yields and rates can then be determined. Many variations of this approach can be used to accommodate the specific requirements of individual enzymes for optimal activity.

[0110] The molecular genetic methods to be used are well-documented. See, for example, Maniatis *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL (Cold Spring Harbor Laboratory, 1989); Ausubel *et al.*, CURRENT PROTOCOLS AND MOLECULAR GENETICS (John Wiley & Sons, 1987). Fermentations can be conducted at the pH levels

described above and in the Examples. Ethanol can be measured by gas-liquid chromatography, as described by Goel *et al.* [1984] *Biotechnol. Lett.* 3:177-182, and Dombek *et al.* [1985] *Appl. Environ. Microbiol.* 51:197-200. Monomeric sugars can be determined colorimetrically and by HPLC. See Ashwell [1966] *Methods Enzymol.* 8:93-95; Wood *et al.* [1988] *Methods Enzymol.* 160. Oligosaccharides can be analyzed by HPLC and thin-layer chromatography. Cultures can be grown in either complex media such as Luria broth or in mineral salts. See Starr, Vol. II THE PROKARYOTES 1166-1172, 1260-1271 (Springer-Verlag, 1981); Mizutani *et al.* [1986] *Biotechnol. Bioeng.* 28: 204-09. Organisms can be maintained on solid medium containing 1.5% agar, and stocks can be stored in glycerol at -70°C.

[0111] Cellulase enzymes can be assayed essentially as described by Curry *et al.* [1988] *Appl. Environ. Microbiol.* 54: 476-484, and by Gripenet *et al.* [1988] *J. Bacteriol.* 170: 4576-4581. Exoglucanase activity can be determined using p-nitrophenylcellobioside as a model substrate. Carboxymethyl cellulase (endoglucanase) activity can be measured as the release of reducing sugar. Cellulase activity also can be assayed by the conventional approach of measuring the release of reducing sugars from filter paper. Methyl-umbelliferyl derivatives, CMC and pectate can be used to prepare indicator plates for qualitative comparisons, thereby to identify strains which are high-level producers of an enzyme hydrolyzing any of these substrates.

[0112] (G) IMPROVING ALCOHOL TOLERANCE AND PRODUCTION OF TRANSFORMANTS: Maximum levels of ethanol produced with ethanologenic recombinants of the present invention can exceed 1 M, for example. Ethanol production can exceed 1.5 M for fermentation of lactose by an *E. coli* transformant, KO11, which is deposited as ATCC 55124. Enhanced alcohol tolerance in the ethanologenic recombinants of the present inventions therefore is particularly useful. Changes affecting membrane lipids and/or stress proteins are likely bases for such improvements. The *Z. mobilis adhB*, a component of a preferred ethanologenic operon (*pet*) within the present invention, also has been identified as a prominent *Z. mobilis* stress protein, responsive both to temperature and to alcohol. See Haejung *et al.* [1991] *J. Bacteriol.* 173: 5975-82. Other *Z. mobilis* genes which encode a stress protein should be useful in the same context. To the end of obtaining such genes, genes such as those found in *E. coli* which express major stress proteins, as reported by Johnson *et al.*, [1989] *J. Bacteriol.* 171: 1590-96, Ang *et al.* [1989] *J. Bacteriol.* 171: 2748-55, Lipinska *et al.* [1988] *Nucleic Acids Res.* 16: 7545-61, and Fayet *et al.* [1986] *Mol. Gen. Genet.* 202: 435-45, can be used as probes to isolate corresponding genes from *Z. mobilis*.

[0113] It is likely that lipid-synthesis genes also will prove useful for ethanol tolerance. Other genes useful for ethanol tolerance can be selected by transforming libraries directly into ethanologens and enriching by serial passages in fermentation broth in which excess sugar levels are maintained to allow further growth of ethanol-resistant organisms after ethanol levels exceed the tolerance of the original inoculum. Chemostat approaches may also be particularly good for these selections.

[0114] Gene libraries of the highly ethanol resistant *Lactobacillus homohiochii* and *L. heterohiochii*, see Ingram [1990] *Critical Rev. Biotechnol.* 9: 305-19, also can be tested in these enrichments. Since these microbes effect spoilage in sake, they tolerate alcohol levels on the order of 15% v/v (over 2.5M). By this approach, therefore, recombinant strains can be developed which should be capable routinely of producing ethanol levels of at least 80 grams per liter.

[0115] Further still, the glycolytic enzymes of *Z. mobilis* may be used to enhance ethanologenic recombinants of the present invention. That is, the expression of one or more of the 13 genes that encode those glycolytic enzymes should increase glycolytic flux and, hence, the rate of ethanol production.

[0116] (H) PROCESS DESIGN: Numerous fermentation processes can be constructed, in accordance with this description, to implement the present invention. Such processes generally will have at least two stages: a "saccharification" stage, where biomass is contacted with polysaccharase to break down the oligosaccharides therein into simpler oligosaccharides and/or monosaccharides, and a subsequent "fermentation" stage in which the simpler oligosaccharides and/or monosaccharides are fermented to ethanol. Advantageously, the two stages will be conducted separately since the optimum conditions for carrying out each stage differ significantly. The optimum conditions for the saccharification stage comprise a temperature of from about 50°C to about 60°C and a pH of from about 4.5 to about 5.0. The optimal conditions for fermentation stage include a temperature of from about 30°C to about 35°C and a pH of about 6.0.

[0117] To enhance the efficiency of the process, there may be some recycling of materials from the fermentation stage to the saccharification stage. Further, since the recycled materials will be at the fermentation stage temperature of from about 30°C to about 35°C, those materials advantageously may be used in a heat exchanger to cool the higher temperature effluent from the saccharification stage before it is introduced into the fermentation stage. An exemplary process is discussed below and illustrated in Figure 19.

[0118] With reference now to Figure 19, polysaccharase and pretreated biomass are provided to the enzyme reactor 10, where the starting oligosaccharides in the biomass are broken down into simpler oligosaccharides and monosaccharides. The enzyme reactor 10 is maintained at a temperature of from about 50°C to about 60°C and a pH of from about 4.5 to about 5.0. A mixture of solids and liquid is drawn from enzyme reactor 10 through conduit 12 and into a solid/liquid separator 14, where the solid and liquid are separated, for example, by centrifugation.

[0119] The separated solids are returned to the enzyme reactor 10 through conduit 16. The effluent from the separator is then passed through conduit 18 to heat exchanger 20, where it is cooled to about 30°C to about 35°C. The effluent

is then passed through conduit 22 to membrane separator 24 which separates large molecular weight polysaccharases, which are then returned via conduit 26 to the enzyme reactor 10. The membrane separator, for example, can be an ultrafilter, such as the CARRE filter by Du Pont®, that has a removal rating of from about 10^{-3} to about 10^{-4} microns. Alternatively, the membrane separator 24 can be eliminated and the effluent introduced directly into fermentor 30. The remaining sugar solution, which comprises oligosaccharides and/or monosaccharides, is then transferred via conduit 28 to the fermentor 30. Prior to entering the fermentor, however, the pH of the sugar solution is adjusted upward, for example, by adding base.

[0120] Carbon dioxide is drawn off from the fermentor through conduit 32. Effluent is drawn from the fermentor through line 34 to a distillation column (not shown) for distillation of ethanol. Mash is drawn from the fermentor and passed through line 36 into a solid liquid separator 40 (e.g., a centrifuge). Microorganisms which are separated in the separator 40 are returned to the fermentor through conduit 42. Effluent from the separator 40 is supplied to heat exchanger 20 to cool the effluent from the solid/liquid separator 14. After leaving the heat exchanger 20, the pH of the effluent is adjusted downward, for example, by adding acid, to from about 4.5 to about 5.0 and then the effluent is introduced to the enzyme reactor 10. Both the enzyme reactor 10 and the fermentor 30 can be provided with mixers 48 and 50, which can be any conventional type.

[0121] Optionally, on a continual basis, fluid (beer) having an ethanol concentration of from about 0.5% to about 5%, and typically from about 1% to about 2%, can be drawn off from fermentor 30 through conduit 52 and fed into evaporator vessel 54, where ethanol and water are evaporated and taken off through line 56 to another distillation stage (not shown). The concentration of ethanol in line 56 can be from about 20% to about 25%. The beer removed from evaporator 54 and returned to the fermentor 30 through line 58 will have an ethanol concentration on the order of less than about 0.5%, i.e., generally about 0.3%. In this way, the ethanol concentration in the fermentor can be maintained at a relatively low level, thereby maintaining a high rate of ethanol production in the fermentor.

[0122] Likewise, the effluent which is returned to the reactor 10 through conduits 36, 44 and 46 will have a lower ethanol concentration and thus provide less inhibition for the reactions in the reactor. Such a continuous process can be driven by the continual addition of biomass to the reactor 10. It may be desirable, however, to discharge the contents of the reactor 10 and the fermentor 30 on a periodic basis in order to discharge built-up solids and lessen the possibility of contamination.

[0123] Thus, the fermentation process discussed above can provide an efficient method of fermenting biomass to ethanol. Those skilled in the art will recognize that many other additional steps and/or stages can be added to the fermentation process. For example, biomass pretreatment stages can be included, in which the biomass for example, is shredded, hydrolyzed (with or without steam and/or acid), treated with polysaccharases and/or other enzymes, etc. Additionally, there can be an initial stage in which recombinant host cells that express and intracellularly accumulate polysaccharases are and heated to lyse the cells and release the polysaccharase. Also, there can be multiple fermentation stages together with additional separations and recycling of solids and effluents along the way.

III. Examples

[0124] The present invention is further described by reference to the following illustrative examples, which should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted. Other methodological aspects of the examples are discussed below.

[0125] Organisms and growth conditions: *Escherichia coli* TC4, see Conway *et al.* [1987a], and plasmid-containing derivatives thereof were used. Plasmids containing the *Z. mobilis* pyruvate decarboxylase gene (pLOI276) and alcohol dehydrogenase B gene (pLOI284) have been described by Conway *et al.* [1987b].

[0126] Strains and growth conditions: Plasmids pUC18 and pUC19 have been described previously, see Yanisch-Perron *et al.* [1985] *Gene* 33: 103-19, as has pLOI204 and pLOI295. See Conway *et al.* [1987b]; Ingram *et al.* [1987]. The construction and properties of pLOI292, pLOI291, pLOI297, pLOI308, pLOI308-2, pLOI308-5 and pLOI308-10 are described below.

[0127] Cultures were grown at 37°C in Luria broth, see Luria & M. Delbruck [1943] *Genetics* 28: 491-511, which contained 50 grams of glucose per liter. Cells for enzyme analyses and inocula for fermentation studies were grown in tubes (13 x 100 mm) containing 3 ml of broth at 37°C in a tube rotator. Overnight cultures were diluted 100-fold into fresh medium. Aerobic cultures (50 ml of broth in 250 ml flasks) were shaken in a reciprocating water bath (120 oscillations per minute). Anaerobic cultures were grown in stoppered serum bottles (100 ml of broth in 130 ml bottles) with gyrator agitation (150 rpm) in a 37°C incubator. Anaerobic cultures were vented with a 25-gauge needle to allow escape of gaseous fermentation products.

[0128] Growth was monitored spectrophotometrically with a Spectronic 70 spectrophotometer (Bausch & Lomb, Inc. Rochester, NY) at 550 nm. Disposable culture tubes (10 x 75 mm) were used as cuvettes. On absorbance unit under our conditions contained approximately 0.25 mg of cellular protein per ml. Growth was measured at A_{550} .

[0129] *Escherichia coli* hosts containing the plasmids of the subject invention have been deposited with the American

Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852 USA. The cultures have been assigned these accession numbers by the repository:

Culture	Accession number	Deposit date
<i>E. coli</i> pLOI308-10	ATCC 67983	May 15, 1989
TC4(pLOI292)	ATCC 68237	February 23, 1990
TC4(pLOI308-11)	ATCC 68238	February 23, 1990
TC4(pLOI297)	ATCC 68239	February 23, 1990
TC4(pLOI295)	ATCC 68240	February 23, 1990
<i>E. coli</i> pLOI510	ATCC 68484	
<i>K. oxytoca</i> M5A1 (pLOI555)	ATCC 68564	March 14, 1991

[0130] The subject cultures have been deposited under conditions that assure that access to the cultures will be available during the pendency of this patent application to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto under 37 CFR 1.14 and 35 USC 122. The deposits are available as required by foreign patent laws in countries where counterparts of the subject application or its progeny are filed. But it should be understood that the availability of the deposits does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

[0131] Further, the subject culture deposits will be stored and made available to the public in accord with the provisions of the Budapest Treaty for the Deposit of Microorganisms. That is, they will be stored with all the care necessary to keep them viable and uncontaminated for a period of at least five years after the most recent request for the furnishing of a sample of a deposit, and in any case, for a period of at least thirty years after the date of deposit or for the enforceable life of any patent which may issue disclosing the cultures. The depositor acknowledges the duty to replace the deposits should the depository be unable to furnish a sample when requested, due to the condition of the deposits. All restrictions on the availability to the public of the subject culture deposits will be irrevocably removed upon the granting of a patent disclosing them.

[0132] Genetic techniques: Transformations, plasmid constructions, DNA digestions and analyses were carried out as previously described. Recombinants were selected on solid media (1.5% agar) containing 2 grams of glucose per liter and appropriate antibiotics. Recombinants containing functional ethanologenic genes from *Z. mobilis* were identified as oversized colonies growing on Luria agar plates which included glucose, and identification was confirmed by the observations of poor growth on Luria agar plates lacking glucose and of alcohol dehydrogenase expression on aldehyde indicator medium.

[0133] Enzyme assays: Cells were disrupted, heat-inactivated, and assayed for pyruvate decarboxylase activity (thermostable) as described previously. See Conway *et al.* [1987b]. Cells were prepared and assayed for alcohol dehydrogenase II activity in the direction of ethanol oxidation as described previously, except that cells were washed and disrupted in 30 mM potassium phosphate buffer to which solid ferrous ammonium sulfate (final concentration, 0.5 mM) and sodium ascorbate (1 mM) had been freshly added. See Neale *et al.* [1986] *Eur. J. Biochem.* 154: 119-24. This modification, combined with the immediate assaying of alcohol dehydrogenase activity without storage, resulted in a much higher specific activity than that previously reported. For the purpose of calculating ethanol-production efficiency, cell mass was measured in terms of protein content per culture, using the Folin phenol reagent method of Lowry *et al.*, *J. Biol. Chem.* 193: 265-75.

[0134] Analysis of fermentation products: Fermentation products were determined in clarified broth with a Millipore/Waters high-performance liquid chromatograph (Millipore Corporation Bedford, MA) equipped with a refractive index monitor and an electronic integrator. Separations were performed on an Aminex HPX-87H column (300 by 7.8 mm) purchased from Bio-Rad Laboratories, Richmond CA, at 65°C at a flow rate of 0.25 ml/min (100 µl injection volume). Peaks were identified by using authentic standards. The two peaks eluting before glucose and the later unknown peak eluting at 45.4 to 45.8 minutes are components of uninoculated medium.

EXAMPLE 1. STRAIN CONSTRUCTION

[0135] The sizes of the structural genes coding for pyruvate decarboxylase and alcohol dehydrogenase II are 1.7 and 1.1 kilobases, respectively, and these genes encode proteins with molecular weights of 60,000 and 40,100. These genes are each located on derivatives of pUC18 under the control of the *lac* promoter (Figure 1). The two genes were combined by inserting the promoterless 1.4 kilobase fragment generated by restriction endonucleases *EcoRI* and *SalI* from pLOI284 (alcohol dehydrogenase) into the *BamHI* site downstream from the pyruvate decarboxylase gene in pLOI276. These clones were selected for resistance to ampicillin and for the presence and expression of alcohol

dehydrogenase activity on a newly developed pararosaniline-ethanol indicator plate which detects the production of aldehydes. Clones containing the indicated construction, pLOI295, grew poorly on the surface of Luria agar plates (aerobic) in the absence of added glucose but grew to much higher densities than the plasmid-free strain and strains containing pLOI276 or pLOI284 on agar plates containing 2% glucose.

[0136] Recombinants containing the *pet* operon were readily detected as larger, more opaque colonies on Luria agar plates (aerobic) containing glucose. This difference in colony size and opacity has proven to be a useful marker for the identification of recombinants which contain plasmids expressing both alcohol dehydrogenase and pyruvate decarboxylase genes.

[0137] The complete base sequence of pLOI295 is known. The open reading frame for the gene coding for pyruvate decarboxylase begins 163 bases downstream from the *lac* promoter and ends with two stop codons 85 bases upstream from the open reading frame of the gene coding for alcohol dehydrogenase II. Both genes include sequences which resemble ribosome-binding sites immediately upstream from each open reading frame. The gene encoding alcohol dehydrogenase II contains a single stop codon followed by a palindromic sequence of 13 base pairs which serves as a transcriptional terminator.

Example 2 - Expression of *Z. mobilis* Genes in *E. coli*

[0138] Both pyruvate decarboxylase and alcohol dehydrogenase II genes were expressed at high levels in *E. coli* under the control of the *lac* promoter singly (pLOI276 and pLOI284, respectively) and jointly. Pyruvate decarboxylase is not present in wild-type *E. coli*, but an inducible alcohol dehydrogenase is present at low concentrations. During growth of *E. coli* in the presence of glucose, the specific activities of the *Z. mobilis* enzymes declined by approximately 50%, which is consistent with glucose repression of the *lac* promoter. The specific activity of pyruvate decarboxylase, coded for by the proximal gene in the *pet* operon, was threefold higher in pLOI295 than in pLOI276. The specific activity of the product of the alcohol dehydrogenase II gene, the distal gene in the *pet* operon, was expressed in pLOI295 at twice the level in pLOI284.

Example 3 - Fermentation of Glucose by Recombinant Strains

[0139] Expression of the *pet* operon in *E. coli* resulted in the production of ethanol as the primary fermentation product during anaerobic growth. The parent strain produced succinate (1.5 mM), lactate (18 mM), and acetate (7 mM) as major fermentation products (Figure 3A). An identical fermentation profile was observed in cells containing pLOI284, which carries the alcohol dehydrogenase II gene (Figure 3C). With pLOI276 carrying the gene coding for pyruvate decarboxylase, an ethanol peak is clearly evident (18 mM), equivalent to one-third of the accumulated fermentation products. This higher level of ethanol results from the combined activities of the pyruvate decarboxylase from *Z. mobilis* and the native *E. coli* alcohol dehydrogenase. With pLOI295 containing the *pet* operon (both pyruvate decarboxylase and alcohol dehydrogenase II genes from *Z. mobilis*), *E. coli* produced large amounts of ethanol (750 mM; 4.3%, vol/vol), which represented over 95% of the fermentation products.

[0140] The high levels of alcohol dehydrogenase and pyruvate decarboxylase produced in cells containing the *pet* operon dominated NADH oxidation in *E. coli*. Thus, the fermentation of this organism was converted to the equivalent of those of *S. cerevisiae* and *Z. mobilis*. During normal fermentative growth, pyruvate is converted to acetyl coenzyme A by the pyruvate dehydrogenase complex, to oxaloacetate (and on to succinate) by phosphoenolpyruvate carboxylase, to formate and acetyl coenzyme by pyruvate formate lyase, and to lactate by lactate dehydrogenase. This last pathway is the dominant route for the regeneration of NAD⁺ in unmodified strains of *E. coli*. But the K_m s for bacterial lactate dehydrogenases are quite high, ranging from 10 to 1,000 mM (Garvie, E.I. [1980] "Bacterial lactate dehydrogenases," Microbiol. Rev. 44:106-139; Tarmy, E.M., and N.O. Kaplan [1968] "Kinetics of *Escherichia coli* B D-lactate dehydrogenase and evidence for pyruvate controlled change in conformation," J. Biol. Chem. 243:2587-2596). The K_m of the pyruvate decarboxylase from *Z. mobilis* is 0.4 mM (Bringer-Meyer, S., K.-L. Schimz, and H. Sahm [1986] "Pyruvate decarboxylase from *Zymomonas mobilis*: Isolation and partial characterization," Arch. Microbiol. 146:105-110). The abundance of this enzyme, coupled with the lower K_m , effectively diverts the flow of pyruvate from glycolysis into ethanol.

[0141] High cell densities are also achieved during mixed growth conditions with moderate agitation or stirring of culture vessels in which gas exchange is not restricted. Under these conditions, a final pH of 6.3 or above was observed, depending upon the extent of aeration.

Example 4 - Plasmid Constructions and Expression of *Z. mobilis* Ethanologenic Enzymes in *E. coli*

[0142] Plasmid pLOI295 contains the *Z. mobilis* genes encoding pyruvate decarboxylase and alcohol dehydrogenase II under the control of the *lac* promoter. This construction is referred to as the *pet* operon and used as the source of

ethanologenic genes for the construction of additional plasmids with alternative promoters. The *EcoRI-SalI* fragment from pLOI295 containing the ethanologenic genes was treated with the Klenow fragment of DNA polymerase to produce blunt ends. This blunt-ended DNA fragment was inserted into the *SmaI* site of pUC19 with the *pdg* gene immediately downstream from the *lac* promoter. The resulting plasmid, denoted pLOI293, contained the *pet* genes flanked by *BamHI* sites. Plasmids pLOI291 and pLOI292 (opposite orientations) were constructed by inserting the *BamHI* fragment containing the genes encoding the ethanologenic enzymes into the expression vector pLOI204. The *BamHI* fragment includes the ribosome-binding site, complete sequences for both genes, and a transcriptional terminator distal to *adhB*. In pLOI292, the two genes are expressed under the control of the *Z. mobilis* promoter contained in the original expression vector.

[0143] Plasmid pLOI308 was constructed to remove the *pet* genes from the control of the *lac* promoter but to retain the upstream *BamHI* site for the insertion of alternative promoters. Partial digestion of pLOI293 with *BamHI* and Klenow treatment were used to remove the *BamHI* site distal to the *adhB* gene. The ethanologenic genes were removed from this plasmid as a promoterless *BamHI* (immediately proximal to *pdg*)-*EcoRI* (distal to *adhB*) fragment, which was directionally inserted into the *BamHI* and *EcoRI* sites of pUC18 to produce pLOI308. This plasmid expressed low levels of *adhB* on aldehyde indicator plates but did not exhibit the large-colony phenotype associated with the other functional *pet* plasmids pLOI295, pLOI291, and pLOI292.

[0144] Chromosomal DNA from *Z. mobilis* was partially digested with *Sau3A* such that most of the DNA appeared to be less than 4 kilobases long. This unfractionated DNA was used as a source of promoter fragments and was ligated into the dephosphorylated *BamHI* site of pLOI308. Ampicillin-resistant recombinants with a well-expressed *pet* operon were identified as large colonies on Luria agar plates containing glucose. Three of the recombinant strains, pLOI308-2, pLOI308-5, and pLOI308-10, were selected for study. The *Z. mobilis* DNA fragments with promoter activity in these plasmids were 6, 2, and 2 kilobases long, respectively.

[0145] Table 1 summarizes the activities of pyruvate decarboxylase and alcohol dehydrogenase in overnight cultures of the recombinant *E. coli*. The activities of pyruvate decarboxylase ranged from 0.37 IU/mg of cell protein in strain TC4(pLOI291) to 8.23 IU in TC4(pLOI295). In terms of pyruvate decarboxylase activity, the recombinant strains of TC4 can be ordered as follows (highest to lowest): pLOI295 > pLOI308-10 > pLOI308-2 > pLOI308-5 > pLOI292 > pLOI291.

Table 1.

Expression of ethanologenic enzymes from <i>Z. mobilis</i> in <i>E. coli</i>				
Plasmid	Pyruvate decarboxylase		Alcohol dehydrogenase	
	Sp act ^a	% Cell protein ^b	Sp act ^a	% Cell protein ^c
pLOI291	0.37	0.4	0.21	0.02
pLOI292	0.48	0.5	0.30	0.03
pLOI308-2	2.26	2.3	1.54	0.21
pLOI308-5	1.11	1.1	0.76	0.10
pLOI308-10	6.5	6.5	2.51	0.34
pLOI295	8.2	8.2	9.65	1.4
None	0	0	0.08	

^a Expressed as micromoles of substrate utilized per minute per milligram of total cellular protein.

^b Calculated assuming a specific activity of 100 for the pure enzyme.

^c Calculated assuming a specific activity of 710 for the pure enzyme after subtraction of native alcohol dehydrogenase activity.

[0146] Alcohol dehydrogenase activities in the recombinant strains followed the same trend in terms of expression from different plasmids as did pyruvate decarboxylase. The alcohol dehydrogenase activities measured represent a combination of the native enzyme from *E. coli* and the *Z. mobilis* enzyme. The level observed in strain TC4 lacking a plasmid was relatively small in comparison to those observed in strains carrying plasmids with the *Z. mobilis* gene. The activities of the *Z. mobilis* enzyme (corrected for native *E. coli* alcohol dehydrogenase) ranged from 0.13 IU/mg of cell protein from strain TC4(pLOI291) to 9.6 IU in TC4(pLOI295).

Example 5 - Growth of Recombinant Strains Containing the Ethanologenic Enzymes From *Z. mobilis*

[0147] Shifting the catabolism of glucose to the production of ethanol also affected growth yield and pH drift of the growth medium. Although fermentation products are relatively nontoxic, they may accumulate to toxic levels during fermentation. During anaerobic growth in bottles containing Luria broth containing 10% glucose, the plasmid-free strain and the strain carrying pLOI284 (carrying the gene coding for alcohol dehydrogenase II) achieved a final density of

0.25 mg of cell protein per ml after 48 hr, with a final pH of 4.4. The cell density increased by twofold in the strain carrying pLOI276 (carrying the gene coding for pyruvate decarboxylase), with a final pH of 4.5. The final cell density of the strain carrying pLOI295 (*pet* operon) was 2.5 mg/ml, 10-fold higher than that of the control strain. The final pH was 4.7. At a density of 2.5 mg of cell protein per ml, magnesium appears to be limiting, and a 1.5-fold further increase in cell density is readily achieved by the addition of 0.5 mM magnesium sulfate.

[0148] The growth of the recombinant strains was examined under both aerobic and anaerobic conditions (Figure 3). Under aerobic conditions (Figures 3A and Table 2), strain TC4 grew with a generation time of approximately 30 min during the most rapid phase of growth. Strain TC4 carrying the derivatives of pLOI308 exhibited equivalent maximal rates of growth, with generation times between 26 and 30 min. Strain TC4(pLOI295) grew poorly under these conditions (generation time, 71 min) and was accompanied by partial lysis. Strains TC4(pLOI291) and TC4(pLOI292) grew at intermediate rates, each with a generation time of 46 min.

Table 2.

Maximal generation times, final cell densities, and final pHs of the broth during aerobic and anaerobic growth.				
Growth pH ^a Condition	Plasmid	Cell density ^a (mg of protein/ml)	Generation time (min)	Final
Aerobic	None	0.7	29	4.4
	pLOI291	0.7	46	5.3
	pLOI292	1.3	46	5.1
	pLOI295	1.1	71	5.7
	pLOI308-2	1.7	27	5.5
	pLOI308-5	0.8	30	5.0
	pLOI308-10	2.5	26	5.0
Anaerobic	None	0.3	32	4.4
	pLOI291	0.4	40	4.5
	pLOI292	1.0	48	5.0
	pLOI295	2.1	39	4.7
	pLOI308-2	0.8	42	5.7
	pLOI308-5	0.4	38	4.9
	pLOI308-10	2.2	41	5.2

^a Measured after 24 hr of growth.

[0149] Under anaerobic conditions (Figure 3B and Table 2), the generation time for strain TC4 lacking a plasmid was 32 min, considerably shorter than that for the recombinant strains containing the ethanologenic enzymes. All of the recombinants exhibited similar maximal rates of growth, with generation times between 38 and 41 min, except for TC4 (pLOI292), which grew somewhat more slowly, with a generation time of 48 min.

[0150] All of the recombinants except TC4(pLOI295) grew after 12 hr under anaerobic and aerobic growth conditions to cell densities equivalent to or higher than those of strain TC4 lacking a plasmid (Figures 3A and B). Table 2 summarizes the final cell densities of strain TC4 and the recombinants after 24 hr of growth. Under aerobic conditions, strain TC4 containing pLOI308-10 reached the highest cell density, followed by TC4 containing pLOI308-2, pLOI292, pLOI295 (with some lysis apparent), and pLOI308-5.

[0151] Under anaerobic conditions, the final cell densities of strain TC4 containing pLOI308-10 and pLOI295 were roughly equivalent, followed by those of TC4 containing pLOI292, pLOI308-2, pLOI308-5, and pLOI291.

[0152] Figure 4 shows the relationship between the level of pyruvate decarboxylase activity in cells and the final cell density after 24 hr of growth. Since the synthesis of pyruvate decarboxylase is coupled to that of alcohol dehydrogenase II in these recombinants, this plot represents the effects of the alternative *Z. mobilis* system for NAD⁺ regeneration on final cell density. From these data, it is clear that the expression of the *Z. mobilis* pathway for the production of ethanol increases final cell density under both anaerobic and aerobic conditions. In strain TC4(pLOI308-10), the levels of expression of pyruvate decarboxylase (6.5 IU) and alcohol dehydrogenase II (2.5 IU) were nearly optimal for both anaerobic and aerobic growth. The level of expression of ethanologenic enzymes in strain TC4(pLOI295) appears to be excessive, resulting in diminished cell growth accompanied by partial lysis under aerobic conditions and slightly reduced growth under anaerobic conditions.

[0153] The increased growth of strain TC4(pLOI295) under anaerobic conditions with little apparent lysis in contrast to the poor growth and lysis during growth in rapidly shaken flasks suggested that a highly aerobic environment may be damaging to this construction. Lysis in this recombinant was dramatically reduced and the final cell density was

increased during growth in shaken flasks when the speed of oscillation was decreased by one-third.

Example 6 - Effects of Ethanologenic Enzymes on the Acidification of Broth During Growth

[0154] Figure 3C shows a plot of the changes in the pH of the broth during anaerobic growth. The pH fell rapidly during the first 6 hr of growth of strain TC4 lacking a plasmid but declined more slowly in derivatives containing the ethanologenic enzymes. Acidification during the initial 12 hr was reduced to the greatest extent in strain TC4 containing pLOI295, followed by TC4 containing pLOI308-10, pLOI308-2, and pLOI308-5. Data for strains TC4(pLOI291) and TC4(pLOI292) are not shown but lie below and above those for TC4(pLOI308-5), respectively. Although the recombinants reached a higher final cell density, the pH of the broth from the recombinants grown under both anaerobic and aerobic conditions for 24 hr was less acidic than that of the broth from strain TC4 lacking ethanologenic enzymes (Table 2).

[0155] The reduced rate and extent of acidification in recombinants accompanied by increased cell growth suggested that the fall in pH was a major factor limiting growth even under highly aerobic conditions. This hypothesis was supported by an 85% increase in the final cell density of strain TC4 (lacking a plasmid) grown in medium supplemented with a 1/10 volume of 1M sodium phosphate buffer (pH 7.0). Lower levels of buffer addition resulted in intermediate levels of growth.

Example 7 - Effects of Ethanologenic Enzymes on Fermentation Products

[0156] Table 3 summarizes the analyses of fermentation products made by strain TC4 and the recombinants after 24 hr of growth under aerobic and anaerobic conditions. Under aerobic conditions, acetate was the primary fermentation product that accumulated during the growth of strain TC4 lacking a plasmid in rich medium, with no detectable ethanol. The amount of acetate produced was drastically reduced in strains containing the ethanologenic enzymes from *Z. mobilis*, and ethanol appeared as the major fermentation product. Strain TC4 containing pLOI308-10 produced the most ethanol, followed by TC4 containing pLOI295, pLOI308-2, pLOI292, pLOI308-5, and pLOI291. Under these aerobic conditions, small amounts of lactate were also produced (0.6 to 1.2 mM) by all of these strains. Only strain TC4 containing pLOI308-10 accumulated appreciable amounts of succinate, although this product still represented only 1% of the total fermentation products, with 94% being ethanol.

Table 3.

Comparison of fermentation products during aerobic and anaerobic growth					
Growth Condition	Plasmid	Fermentation product [mM (SD)]			
		Succinate	Lactate	Acetate	Ethanol
Aerobic	None	0.2 (0.1)	0.6 (0.2)	55 (2)	Tr
	pLOI308-2	Tr	1.2 (0.3)	22 (2)	98 (3)
	pLOI308-5	Tr	0.9 (0.2)	43 (3)	15 (2)
	pLOI308-10	4.9 (0.5)	1.0 (0.2)	17 (2)	337(21)
	pLOI295	Tr	1.1 (0.4)	13 (1)	114(10)
	pLOI291	Tr	0.6 (0.2)	34 (3)	7 (1)
	pLOI292	Tr	1.3 (0.2)	30 (1.5)	24 (1)
Anaerobic	None	0.9 (0.1)	22 (1)	7 (0.3)	0.4 (0.2)
	pLOI308-2	0.8 (0.1)	7 (0.5)	4 (0.3)	71 (5)
	pLOI308-5	0.3 (0.1)	18 (2)	6 (1)	16 (2)
	pLOI308-10	5.0 (0.4)	10 (1)	1.2 (0.2)	482 (23)
	pLOI295	2.2 (0.20)	6 (1)	3 (0.3)	90 (2)
	pLOI291	1.0 (0.1)	15 (0.5)	7 (0.2)	4 (0.5)
	pLOI292	2.3 (0.2)	9 (0.7)	7.2 (0.3)	21 (1)

[0157] Under anaerobic conditions, lactate was the principal fermentation product that accumulated during 24 hr of growth of strain TC4 lacking a plasmid in rich medium containing glucose, with lesser amounts of acetate, succinate, and ethanol being present. Lactate production was dramatically reduced in strains containing the ethanologenic enzymes and was accompanied by the production of substantial quantities of ethanol. Strain TC4(pLOI308-10) produced

the largest amount of ethanol, and this product alone represented 97% of the total soluble fermentation products. The trend of ethanol production among the organisms tested was the same as that during aerobic growth. All organisms except TC4(pLOI308-10) actually produced less total ethanol after 24 hr under anaerobic conditions than under aerobic conditions. It is likely that this low level of accumulated ethanol was caused by the reduction in total cell mass produced under these anaerobic conditions, thus reducing the volumetric rate of ethanol production.

[0158] The extent of ethanol production under anaerobic and aerobic conditions (Table 3) was directly related to the level of expression of the *Z. mobilis* ethanologenic genes (Table 1). Ethanol production appeared to be optimal in strain TC4(pLOI308-10), with a pyruvate decarboxylase activity of 6 IU and alcohol dehydrogenase II activity of 2.5 IU.

[0159] Derivatives of *E. coli* TC4 containing plasmids which express the ethanologenic enzymes from *Z. mobilis* grew to higher cell densities than did the parent organism lacking a plasmid. The increase in the final cell density, the extent to which ethanol accumulated in the medium, and the reduction in the rate of acidification of the culture broth during growth all correlated with the level of expression of *Z. mobilis* ethanologenic enzymes. Heterologous promoters were used to express the genes in all constructions except pLOI295 (*lac*) to minimize potential problems associated with transcriptional regulation. The level of expression nearest to optimal for growth and ethanol production was provided by pLOI308-10 (6.5 IU of pyruvate decarboxylase and 2.5 IU of alcohol dehydrogenase II per mg of total cellular protein). This level of expression in *E. coli* is considerably higher than that present in *Z. mobilis* CP4, which contains only the ethanol pathway for the regeneration of NAD⁺.

[0160] The level of expression of ethanologenic enzymes appeared to be excessive in strain TC4(pLOI295) (approximately 17% of the soluble cellular protein). This high level of expression was accompanied by partial cell lysis, slower growth, and a reduction in ethanol production under aerobic conditions. These effects were reduced by slower agitation and by growth under anaerobic conditions. The apparent damage and partial lysis that occurred during highly aerobic growth may have been related to the depletion of NADH by a combination of the high levels of *Z. mobilis* alcohol dehydrogenase II and the native NADH oxidase (coupled to the electron transport system).

[0161] The production of ethanol as a major product does not appear to adversely affect the growth rate of *E. coli* TC4. Strains containing derivatives of pLOI308 (ColE1 replicon) expressing the *pet* operon and producing ethanol grew as rapidly as did the parent organism under aerobic conditions with glucose and reached higher final cell densities than did the parent organism. Strains containing pLOI291 or pLOI292 with the RSF1010 replicon grew more slowly under aerobic conditions. Since these two constructions expressed lower levels of the ethanologenic enzymes and produced less ethanol than did pLOI308-10, the reasons for the slower growth can be attributed to properties of the vector rather than to the expression of the *pet* operon.

EXAMPLE 8. PREPARATION OF ADDITIONAL STRAINS

[0162] Additional *E. coli* strains were tested in order to identify bacteria with superior characteristics for use as an ethanol producing microbe. The following *E. coli* strains were evaluated: ATCC 8677, ATCC 8739, ATCC 9637, ATCC 11303, ATCC 11775, ATCC 14948, ATCC 15244, and ATCC 23227. These were grown in a shaking water bath at 30°C in Luria broth (Luria, S.E. and M. Delbruck [1943] Genetics 28:491-511) containing tryptone (10 g/liter), yeast extract (5 g/liter), sodium chloride (5 g/liter), and a fermentable sugar. Glucose and lactose were added at concentrations of 100 g/liter and xylose at a concentration of 80 g/liter unless indicated otherwise. Sugars were autoclaved separately (121°C, 15 min), double strength in distilled water. Xylose (Sigma Chemical Co., St. Louis, MO) solutions were acidic and were neutralized with sodium hydroxide prior to autoclaving; failure to neutralize resulted in extensive browning and decomposition. Similar fermentation results were obtained with sugars which were autoclaved or filter-sterilized. Survival in broth and on plates of recombinant strains containing genes encoding the enzymes of the ethanol pathway required the presence of a fermentable sugar. Where indicated, tetracycline was added at a final concentration of 10 mg/liter.

Example 9 - Environmental Hardiness of Additional Strains

[0163] Prior to the introduction of plasmids for ethanol production, the growth of the previously-identified 8 different strains of *E. coli* were compared for environmental hardiness. Strains were tested for their resistance to sodium chloride, sugars, low pH, and ethanol. Concentrations of sodium chloride and sugars in Table 4 include that present in the original medium. The original pH of the medium was 6.8; this was adjusted to lower values with HCl where indicated. Acidified media were sterilized by filtration. Ethanol was added to autoclaved medium after cooling. Sugars were autoclaved separately. Overnight cultures grown in each respective sugar in the absence of test agent were diluted 60-fold into 13 x 75 mm culture tubes containing 3 ml of test media. Growth was measured as O.D. at 550 nm after 48 hours. An O.D. of 1.0 is equivalent to 0.25 mg/ml of cell protein and 0.33 mg of cell dry weight. In tests of environmental hardiness, a final O.D. below 0.2 reflected less than two doublings and was considered negligible.

[0164] Table 4 summarizes these results in medium containing glucose. Similar results were obtained in medium

EP 0 576 621 B1

containing lactose or xylose. Strains ATCC 8677, ATCC 8739, and ATCC 11775 were particularly sensitive to inhibition by low concentrations of ethanol. Strains ATCC 9637 and ATCC 11775 were the most resistant to low pH although all strains grew for at least 2 to 4 doublings at pH 4.0 except ATCC 23227. All strains grew at 45°C with limited growth at higher temperatures; none could be subcultured above 45°C. All strains grew in media containing 20% glucose, 20% lactose, or 12% xylose.

Table 4.

Growth of <i>E. coli</i> in Luria broth containing 100 g/l glucose under chemical and physical stresses.								
Stress	Growth of <i>E. coli</i> strains (ATCC numbers)							
	8677	8739	9637	11303	11775	14948	15224	23227
NaCl (g/l)								
50	+	++	++	++	++	++	++	++
60	0	+	++	++	+	++	+	++
70	0	0	+	+	0	+	+	+
Ethanol (% by vol.)								
3.8	++	++	++	++	++	++	++	++
5.0	+	+	++	+	+	+	+	+
6.3	0	++	+	+	+	+	+	0
7.5	0	+	+	0	0	+	0	0
8.8	0	0	0	0	0	0	0	0
Acidity								
pH 4.50	++	++	++	++	++	++	++	++
pH 4.25	++	++	++	+	++	++	++	+
pH 4.00	+	+	++	+	++	+	+	0
pH 3.75	0	0	+	0	+	0	0	0
Temperature (°C)								
45	++	++	++	++	+	++	++	++
47	+	+	+	+	+	+	+	+
49	0	0	+	+	0	0	+	+
0 = less than two doublings in O.D. 550 nm								
+ = two to four doublings								
++ = over four doublings								

Example 10 - Sugar Utilization of Additional Strains

[0165] Sugar utilization was tested in two ways. Strains which developed red colonies on MacConkey agar supplemented with 2% carbohydrate were scored positive for sugar utilization (Silhavy, T.J. and J.R. Beckwith [1985] Microbiol. Rev. 49:398-418). Cells were also tested using the Biolog EC plates (Biolog, Inc., Hayward, CA) according to the directions of the manufacturer. The Biolog plates were rapid and convenient, detecting NADH production (conversion of a tetrazolium salt to the insoluble formazan) as a measure of substrate utilization. Both methods were in complete agreement for the 13 sugars examined.

[0166] All strains tested utilized glucose, fructose, galactose, mannose, arabinose, lactose, glucuronic acid, and galacturonic acid. Strain 11775 did not utilize xylose. Maltose and maltotriose were not used by ATCC 11303 and ATCC 23227. All strains exhibited a weak positive reaction with cellobiose. Only strain ATCC 9637 utilized sucrose. The results of the sugar utilization studies are shown in Table 5.

Table 5.

Growth of <i>E. coli</i> strains harboring pLOI297 and pLOI308-11.									
Sugar	Plasmid	Final O.D. 550 nm of <i>E. coli</i> strains (ATCC numbers)							
		8677	8739	9637	11303	11775	14948	15224	23227
Glucose	none	4.0	3.7	6.1	6.0	4.7	5.6	7.0	6.6
Glucose	pLOI297	10.0	10.5	10.5	10.0	9.5	-	9.5	10.2
Glucose	pLOI308-11	9.8	9.5	11.4	11.2	-	9.3	10.8	11.4
Lactose	none	4.3	3.8	7.5	6.0	4.5	6.1	7.0	6.4
Lactose	pLOI297	13.0	6.8	11.6	10.8	7.6	-	10.5	7.0
Lactose	pLOI308-11	10.0	10.0	11.5	11.0	-	7.3	10.0	10.0
Xylose	none	4.1	3.7	7.7	7.3	4.9	5.9	7.2	7.0
Xylose	pLOI297	8.1	10.6	10.8	10.6	4.7	-	11.0	11.0
Xylose	pLOI308-11	10.0	6.8	11.4	8.5	-	11.4	10.6	12.0

Dashed lines indicate no data available.

Example 11 - Genetic Alteration of Additional Strains

[0167] Two new plasmids were constructed using standard methods (Maniatis, T., E.F. Fritsch, and J. Sambrook [1982] Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) which contained resistance genes for ampicillin and tetracycline as selectable markers. The ethanol production operon (*pet*-operon) containing a cryptic *Z. mobilis* promoter, pyruvate decarboxylase, alcohol dehydrogenase and transcriptional terminator was removed as a 5.2 kb *EcoRI* fragment from pLOI308-10 (Ingram and Conway [1988], *supra*) and inserted into the *EcoRI* site of pBR322 to produce pLOI308-11. The plasmid pLOI297 was constructed by inserting the 2.6 kb *EcoRI* fragment from pCOS2EMBL (Poustka, A., H.R. Rackwitz, A.-M. Firschauf, and H. Lehrach [1984] Proc. Natl. Acad. Sci. USA 81:4129-4133) containing the tetracycline resistance gene in to the *SalI* site of pLOI295 (Ingram et al. [1987], *supra*). Cohesive ends were removed by treatment with the Klenow fragment of *E. coli* DNA polymerase (Maniatis et al., *supra*), prior to the ligation.

[0168] Plasmids were introduced into the different strains of *E. coli* by transformation using the calcium chloride procedure of Mandel and Higa (Mandel, M., and A. Higa [1970] J. Mol. Biol. 53:159-162). Selections were made on solid medium containing 2% glucose and tetracycline. Plasmid stability is expressed as the percentage of cells retaining antibiotic markers after 25 generations of growth in the absence of antibiotic selection.

[0169] Recombinant strains harboring plasmids with the genes for ethanol production grew as unusually large colonies which became yellow after 24 to 48 hours on solid medium containing a fermentable sugar. In liquid medium, the final cell densities of these recombinants were twice to three times higher than that of the control lacking plasmid. No transformants were obtained after several attempts from ATCC 14948 with pLOI297 or from ATCC 11775 with pLOI308-11. Strain ATCC 11775 did not utilize xylose and recombinants of this strain did not grow to higher densities than the control with xylose as the fermentable sugar, although increased growth was observed with lactose and glucose.

[0170] Plasmid stability was examined after growth in medium containing glucose for 25 generations (Table 6). Both plasmids contained the same replicons and were maintained well in all strains except ATCC 8677 and ATCC 8739.

Table 6.

Stability of pLOI297 and pLOI308-11 after 25 generations of growth with glucose in the absence of antibiotic selection.		
ATCC Strain	Cells retaining plasmid (%)	
	pLOI297	pLOI308-11
8677	75	85
8739	44	47
9637	100	90
11303	98	98
11775	100	---
14948	---	97

Table 6. (continued)

Stability of pLOI297 and pLOI308-11 after 25 generations of growth with glucose in the absence of antibiotic selection.		
ATCC Strain	Cells retaining plasmid (%)	
	pLOI297	pLOI308-11
15224	99	100
23227	91	100
Dashed lines indicate no data available.		

Example 12 - Expression of Pyruvate Decarboxylase Activity in Genetically Altered Strains

[0171] Pyruvate decarboxylase activity was measured as previously described (Conway et al. [1987], *supra*; Neale et al. [1987], *supra*) except that cells were harvested at an O.D. of 4.0, approximately half maximal growth.

[0172] The expression of *Z. mobilis* pyruvate decarboxylase activity was examined after growth in the presence of tetracycline (Table 7. With pLOI297, *Z. mobilis* genes are expressed under the control of the *E. coli* lac promoter; pLOI308-11 utilizes a cryptic *Z. mobilis* promoter for expression of *pet*-operon. Strains ATCC 11303(pLOI297), ATCC 11775(pLOI297) and ATCC 15224(pLOI297) contained the highest levels of activity.

Table 7.

Expression of <i>Z. mobilis</i> pyruvate decarboxylase in <i>E. coli</i> strains harboring pLOI297 and pLOI308-11 during growth with glucose.		
ATCC Strain	Pyruvate decarboxylase activity ^a	
	pLOI297	pLOI308-11
8677	5.7	6.0
8739	0.8	1.4
9637	1.1	1.4
11303	16.7	2.1
11775	17.1	— ^b
14948	— ^b	2.5
15224	16.3	1.8
23227	2.3	1.7

^a Activity in I.U./mg cell protein.

^b Dashed lines indicate no data available.

Example 13 - Ethanol Production by Genetically Altered Strains

[0173] Luria broth was modified for fermentation experiments by the inclusion of potassium phosphate buffer (pH 7.0) at a final concentration of 0.2 M. Phosphate buffer, complex medium components, and sugars were autoclaved separately and mixed after cooling. Tetracycline was included at a concentration of 10 mg/liter. Inocula were grown from freshly isolated colonies for 24 hours, washed in the fermentation medium to be tested, and added to an initial O.D. 550 nm of approximately 1.0. Fermentations were carried out at 30°C or 37°C in 100 ml volumetric flasks containing 80 ml of broth, fitted with rubber septa and 25 gauge needles to allow gas escape. Fermentation flasks were immersed in a temperature-controlled water bath and stirred by a magnetic stirrer at 100 rpm.

[0174] Ethanol concentration was measured by gas chromatography as previously described (Dombek, K.M. and L. O. Ingram [1985] Appl. Environ. Microbiol. 51:197-200) and is expressed as percentage by volume. The conversion efficiency was calculated based upon sugar added, assuming that 100% efficiency results in the production of 12.75 ml of ethanol (10.2 g) per 20 g of glucose or xylose and 13.5 ml of ethanol (10.8 g) per 20 g of lactose.

[0175] All genetically engineered strains of *E. coli* produced significant amounts of ethanol from sugars (Table 8). Preliminary experiments with strain ATCC 15244(pLOI297) indicated that higher levels of ethanol were produced in medium containing 0.2 M potassium phosphate buffer (pH 7.0). It is anticipated that similar or superior results would be obtained using an automated pH adjustment in place of this buffer. With 15% glucose, higher ethanol levels were produced at 30°C than at 37°C after 48 hours. The fermentation of lactose and xylose were examined only at the lower temperature, 30°C. In general, higher levels of ethanol were produced by strains harboring pLOI297 than with

pLOI308-11. Strains ATCC 11303(pLOI297), ATCC 11775(pLOI297) and ATCC 15224(pLOI297) produced the highest levels of ethanol after 48 hours from 15% glucose, 5.8% to 6.9% by volume. Most strains were less tolerant of xylos in initial experiments and comparisons of fermentation were carried out using 8% xylose. Strains ATCC 9637(pLOI297), ATCC 11303(pLOI297), and ATCC 15224(pLOI297) produced the highest lev ls of ethanol (4.8% to 5.2%) from 8% xylos after 72 hours. All strains grew well in 15% lactose. Strains ATCC 11303(pLOI297) and ATCC 15224(pLOI297) produced the highest levels of ethanol from lactose after 48 hours, 6.1% and 5.6%, respectively.

Table 8.

Ethanol produced in batch fermentation from glucose (48 hours), xylose (72 hours), and lactose (48 hours) by <i>E. coli</i> strains harboring pLOI297 and pLOI308-11.									
Carbohydrate	Plasmid	Ethanol (% v/v) produced by <i>E. coli</i> strains (ATCC numbers)							
		8677	8739	9637	11303	11775	14948	15224	23227
15% glucose	pLOI297 ^a	2.4	4.7	4.2	4.3	4.8	-	4.8	0.9
15% glucose	pLOI308-11 ^a	3.6	1.4	2.1	1.3	-	3.4	2.8	1.3
15% glucose	pLOI297 ^b	3.2	4.7	4.1	5.8	6.9	-	6.1	3.1
15% glucose	pLOI308-11 ^b	5.8	5.0	3.5	1.5	-	3.8	3.0	3.2
15% lactose	pLOI297 ^b	2.3	4.4	5.3	6.1	4.5	-	5.6	3.7
15% lactose	pLOI308-11 ^b	2.3	2.1	3.4	0.9	-	2.9	2.7	3.0
8% xylose	pLOI297 ^b	0.9	4.1	4.8	5.2	-	-	4.8	1.2
8% xylose	pLOI308-11 ^b	2.0	2.8	2.8	1.2	-	2.0	3.5	1.0
Dashed lines indicate no data available.									

^a Incubated at 37°C.

^b Incubated at 30°C.

[0176] Based upon these comparative studies, strains ATCC 11303(pLOI297) and ATCC 15224(pLOI297) appeared to be the best constructs for ethanol production. The time course of growth and ethanol production were examined with both strains in 12% glucose, 12% lactose, and 8% xylose (Figure 1). Cell mass increased approximately 10-fold, reaching a final density of 3.6 g dry weight/liter. With xylose, cell mass increased at half the rate observed with glucose or lactose. Ethanol production and growth were approximately linear for the three sugars until the concentration of ethanol reached 5%.

[0177] To compute the conversion efficiency of sugar to ethanol, final ethanol concentrations after 120 hours were averaged from three sets of experiments (Table 6). The final concentration of ethanol in cultures grown with 12% glucose was 7.2% (by vol.), representing 94% of theoretical yield from glucose. With 12% lactose, the final ethanol concentration was 6.5%, 80% of the theoretical yield from lactose. With 8% xylose, we consistently obtained yields of 100% and higher. These high yields during slower growth with xylose may reflect the conversion of pyruvate from the catabolism of complex nutrients into ethanol, in addition to pyruvate from glucose.

[0178] The rate of ethanol production was computed from the graphs in Figure 1 and are summarized in Table 9. Volumetric productivity of ethanol ranged from 1.4 g/liter per hour for glucose to 0.64 g/liter per hour for xylose. Specific productivity of ethanol was highest during the initial growth period for each of the three sugars. The highest productivity was obtained with glucose, 2.1 g ethanol/g cell dry weight per hour. The highest yield of ethanol per g of sugar was obtained with xylose, exceeding the maximal theoretical yield for sugar alone.

Table 9.

Averaged kinetic parameters for ethanol production by ATCC 11303(pLOI297) and ATCC 15224(pLOI297).					
Sugar	Volumetric ^a Productivity	Specific ^b Productivity	Yield ^c	Efficiency ^d	Ethanol ^e
12% glucose	1.4	2.1	0.48	95%	58
12% lactose	1.3	2.0	0.43	80%	52

^a g ethanol/g sugar

^b ethanol produced/theoretical maximum from sugar substrate x 100

^c ethanol produced/theoretical maximum from sugar substrate x 100

^d final ethanol concentration in g/liter

^e ethanol produced/theoretical maximum from sugar substrate x 100

Table 9. (continued)

Averaged kinetic parameters for ethanol production by ATCC 11303(pLOI297) and ATCC 15224(pLOI297).					
Sugar	Volumetric ^a Productivity	Specific ^b Productivity	Yield ^c	Efficiency ^d	Ethanol ^e
8% xylose	0.6	1.3	0.52	102%	42

^a g ethanol/liter per hour^b g ethanol/g cell dry weight per hour^c g ethanol/g sugar^d ethanol produced/theoretical maximum from sugar substrate x 100^e final ethanol concentration in g/liter

[0179] Experiments were conducted with ATCC 11303(pLOI297) to examine ethanol production from arabinose, galactose, and mannose. Ethanol concentrations of 3% to 4% were obtained after 48 hours at 30°C but were not investigated further. These sugars were metabolized by pathways similar to those for glucose and xylose and would be expected to produce analogous yields (Lin, E.C.C. [1987] "Dissimilatory pathways for sugars, polyols, and carboxylates," In F.C. Neidhardt, J.L. Ingraham, K.B. Low, B. Magasanik, and M. Schaechter [eds], *Escherichia coli* and *Salmonella typhimurium*, Vol. 1, pp. 244-284. American Society for Microbiology, Washington, D.C.).

Example 14 - Ethanol Production from Hosts Other than *E. coli*

[0180] The great similarity in pathways between various organisms makes it predictable that a variety of hosts can be transformed with genes which then confer upon the transformed host the ability to produce ethanol as a fermentation product. When hosts are transformed with the *adh* and *pdc* genes, as described herein, it is fully predictable that appropriate expression vectors can be utilized to achieve expression. Once expression is achieved, the production of ethanol is highly predictable because of the uniformity of this pathway across various potential hosts.

Example 15 - Ethanol Production in Transformed *Klebsiella*

[0181] The *Klebsiella* strain used in this study was previously designated as *Klebsiella pneumoniae* strain M5A1 (Mahl, M.C., P.W. Wilson, M.A. Fife, W.H. Ewing [1965] J. Bacteriol. 89:1482-1487). This strain, a dinitrogen-fixing organism, was originally identified as *Aerobacter aerogenes* but renamed based upon antigenic properties. Since new taxonomic criteria for *Klebsiella pneumoniae* were defined in the 8th edition of Bergy's Manual, the speciation of this new strain was investigated further. Strain M5A1 grew at 10 C in glucose minimal medium but failed to produce gas from lactose at 44.5 C. This strain was indole positive and utilized both m-hydroxybenzoate and gentisate as sole sources of carbon for growth at both 30 C and 37 C. Based upon these tests, M5A1 was designated *K. oxytoca*. Strains were subcultured on Luria agar plates lacking added sugar unless they harbored plasmids encoding *Z. mobilis* genes. Recombinants containing *adhB* and *pdc* require a fermentable carbohydrate for survival and were maintained on plates containing 2% glucose or xylose. Antibiotic concentrations were as follows: ampicillin, 50 µg/ml; chloramphenicol, 40 µg/ml; and tetracycline, 12.5 µg/ml. Expression of *Z. mobilis adhII* in recombinants was screened using aldehyde indicator plates. *Escherichia coli* strain TC4 was used as the host for all plasmid constructions.

[0182] Since M5A1 is relatively resistant to penicillin and its derivatives, we have constructed *E. coli* shuttle vectors carrying the *cat* (Cm^r) or *tet* (Tc^r) genes. A *tet* gene was added to pLOI276 containing *Z. mobilis pdc* by inserting a 2.6-kilobase pair *EcoRI* fragment from pcos2EMBL (Poustka, A., H.R. Rackwitz, A.-M. Frischauf, B. Hohn, H. Lehrach [1984] Proc. Natl. Acad. Sci. USA 81:4129-4133) into the *SalI* site of pLOI276. Cohesive ends were removed by treatment with the Klenow fragment of *E. coli* DNA polymerase before ligation. The resulting construct was confirmed by restriction analysis and designated pLOI560.

[0183] Preliminary studies indicated that *E. coli* B (ATCC 11303) harbored cryptic low copy number plasmids. A new and useful vector was constructed *in vivo* by integrating the *Z. mobilis pdc* and *adhB* along with *cat* into these plasmids. This was done by isolating a 4.6-kilobase pair, promoterless *PstI* fragment containing *pdc*, *adhB*, and *cat* from pLOI510. No replication functions are present on this fragment. After circularizing by self-ligation, these fragments were transformed into *E. coli* B with selection for Cm^r on Luria agar plates containing 2% glucose. Transformants were tested on aldehyde indicator plates and dark red clones were selected for high level expression of the *adhB* gene. Plasmid preparations from these strains were tested for their ability to transfer antibiotic resistance and *Z. mobilis* genes into *E. coli* TC4 by transformation. All recombinants were sensitive to ampicillin, indicating a lack of the pUC18 fragment containing *bla* and the *colE1* replicon. One of these, pLOI555 (8.4 kilobase pair), produced the most intensely red colonies on aldehyde indicator plates, conferred excellent ethanol production ability to *E. coli*, and appeared to be

present in low copy number based upon yields from small scale plasmid isolations. This plasmid, as well as pLOI297 and pLOI560, were used to transform *K. oxytoca* M5A1 with a selection for *Cm^r*.

[0184] Three plasmid constructs containing the *Z. mobilis* gene encoding PDC were transformed into M5A1 (Table 10). The levels of PDC were 8-fold higher in the two pUC-based constructs (pLOI297 and pLOI560) than with pLOI555. Assuming a maximum specific activity for pure PDC of 100 U, this enzyme comprises over 25% of the cytoplasmic protein in M5A1(pLOI297) and M5A1(pLOI560) and 3.6% in M5A1(pLOI555).

Table 10.

Plasmid stability and <i>pdc</i> expression in recombinant strains of M5A1			
Plasmid	PDC activity after 24 hours (U/mg protein)	Percentage retaining traits ^{a,b} (number of generations)	
pLOI555	3.6	100(38.5)	98(68.5)
pLOI297	28	52(38.6)	10(68.6)
pLOI560	27	97(32.9)	0(62.9)

^a Cells were sampled at two times during cultivation.

^b Both the aldehyde trait and antibiotic resistance were lost concurrently.

[0185] The expression of *Z. mobilis* genes in M5A1 was further confirmed in SDS-PAGE gels. Bands which contain PDC and ADHII were easily identified by comparison with the native strain. The band containing PDC is much larger in pUC-based recombinants than in M5A1(pLOI555), consistent with measurements of enzymatic activity. Although ADHII is less abundant than PDC, relative expression of this *Z. mobilis* gene is also higher in M5A1(pLOI297) than in M5A1(pLOI555). No band corresponding to ADHII was evident in M5A1(pLOI560) which contains only the *Z. mobilis* *pdc* gene.

[0186] The copy number of pLOI555 was estimated to be less than 1/10 that of the two other constructs based on yields in small scale plasmid preparations. Although this estimate is only approximate, it is clear that the high levels of PDC present in the pUC-based constructs are due in part to higher copy number.

[0187] Cells harboring pLOI555, pLOI297, or pLOI560 were serially transferred in Luria broth containing 10% glucose without antibiotics for more than 60 generations at 30 °C. Appropriate dilutions of cultures were plated on Luria agar containing 2% glucose without antibiotics. Colonies were tested on aldehyde indicator plates for the retention of the ethanol production genes from *Z. mobilis* and for resistance to appropriate antibiotics.

[0188] * Excessive instability of *Z. mobilis* genes in pBR322-based vectors was reported previously for *Klebsiella planticola* (Tolan, J.S., R.K. Finn [1987] Appl. Environ. Microbiol. 53:2039-2044) and would not be acceptable for industrial processes. Recombinants harboring pLOI555 were very stable with 98% of the population retaining both the antibiotic resistance gene and the genes from *Z. mobilis*. Although only *adhB* expression is detected by aldehyde indicator plates, these recombinants also retained the large colony phenotype indicative of expression of both *pdc* and *adhB*.

[0189] Fermentations were carried out in Luria broth containing 10% (w/v) glucose or xylose at 30 °C, pH 6.0, 100 rpm agitation. Inocula were grown overnight at 30 °C from isolated colonies in unshaken flasks. Fermentations were inoculated to an initial O.D._{550nm} of 1.0 (330 mg dry weight of cells/liter). A Bausch and Lomb Spectronic 70 spectrophotometer was used to monitor growth.

[0190] Ethanol concentrations were determined by gas-liquid chromatography. Conversion efficiencies were corrected for volume changes caused by the addition of base and assumed that all sugars had been metabolized. The maximum theoretical yield of ethanol from xylose and glucose was calculated to be 0.51 g ethanol per gram of sugar with the balance as carbon dioxide. Volumetric productivities were estimated from the most active periods and represent maximum values. All fermentation data in tables and in figures are averages from two or more batch fermentations.

[0191] Table 11 shows the effects of *Z. mobilis* PDC+ADHII on ethanol production from glucose and xylose respectively to the native strain M5A1. M5A1(pLOI555) was clearly the best construct for ethanol production with maximum volumetric productivities of 2.1 g/liter per hour for both glucose and xylose. With either sugar, this recombinant produced approximately 37 g ethanol/liter after 30 hours. Fermentation of these sugars was essentially completed after 48 hours with 45 g ethanol/liter.

Table 11. Ethanol production from glucose and xylose by recombinant strains of M5A1^a

Plasmid	Genes ^b	Base (mmol/g of sugar) ^c	Time ^d (hr)	Ethanol yield g/l g/g of sugar	Theoretical yield (%)	VP (g/liter per hour) ^e	30-h ethanol (g/liter)	Cell yield (g/g sugar)
Glucose								
no plasmid		1.0	48	15	31	1.1	15	0.044
pLOI560 <i>pdg</i>		0.8	72	44	90	1.1	15	0.018
pLOI297 <i>pdg adhB</i>		0.7	72	50	102	1.3	24	0.020
pLOI555 <i>pdg adhB</i>		1.0	48	48	98	2.1	43	0.040
Xylose								
no plasmid		1.9	96	14	31	0.5	7	0.044
pLOI560 <i>pdg</i>		0.7	96	37	75	1.2	5	0.025
pLOI297 <i>pdg adhB</i>		0.9	96	37	76	1.0	12	0.024
pLOI555 <i>pdg adhB</i>		1.1	48	46	94	2.0	37	0.054

^a Calculations are based on total sugar added initially.^b Genes indicated are from *Z. mobilis*.^c Amount of base consumed to maintain a pH of 6.0 during fermentation.^d Time of maximum ethanol concentration.^e VP, maximum volumetric productivity during batch fermentation.

[0192] As with ethanol production, the growth of M5A1(pLOI555) was clearly superior. Growth of this recombinant

was almost equivalent to that of the parental strain. However, unlike the parental strain, cell density progressively declined after reaching a maximum at 15 hours. This decline may reflect a reduction in refractility as ethanol accumulated since lysis was not evident. Without the addition of base to control pH, recombinants containing *pdc+adhB* grew to over twice the density of the parent organism due to a reduced rate of acid production (higher proportion of neutral fermentation products) as observed previously with *E. coli*.

[0193] The maximal volumetric productivities (2.1 g ethanol/liter per hour) are almost double that of *E. coli* recombinants while maintaining similarly high efficiencies and final ethanol concentrations. Unlike *E. coli*, M5A1 (pLOI555) ferments xylose and glucose at equivalent rates. Plasmid pLOI555 was stably maintained in M5A1 in the absence of antibiotic selection. Since the range of substrates for M5A1 is equivalent to that of *E. coli*, M5A1 recombinants offer a distinct and unexpected advantage for ethanol production.

Example 16 - Fermentation of Cellobiose to Ethanol by ethanologenic recombinants of *Erwinia* and of *Klebsiella*

[0194] This example illustrates the fermentation of an oligosaccharide, cellobiose, using ethanologenic recombinants of *Erwinia* and *Klebsiella*.

[0195] The *Erwinia* used in this example was a typical strain of *Erwinia carotovora*. The *Erwinia* was transformed using the plasmid pLOI555 described above in Example 15. The plasmid pLOI555 was inserted into the *Erwinia* using a Biorad Electroporation unit and following the procedures of Ito *et al.* [1988] *Ag. and Biol. Chem.* 52(1): 293-4 (voltage = 2.0 kV, capacitance = 25 μ F, and resistance = 100 ohms). High numbers of transformants, i.e., more than 100,000 per μ g of plasmid DNA, were recovered.

[0196] The *Klebsiella* used in this example was *Klebsiella oxytoca* M5A1 strain P2, which was obtained as described below in Example 18.

[0197] The fermentations were conducted as in Example 15, using 10% cellobiose in Luria broth (pH 6, 30°C) and 100 rpm agitation. The concentration of ethanol produced by the *Erwinia* and *Klebsiella oxytoca* was 1.113 molar and 1.065 molar, respectively. The results are provided in Figure 6 in g/L of ethanol produced. Symbols: ■, *Erwinia*; □, *Klebsiella*.

Example 17 - Fermentation of Polymeric Feedstocks to Ethanol by a Single, Genetically Engineered Microorganism

[0198] **Summary of Example:** This example illustrates a two-stage process for fermenting polymeric feedstocks to ethanol by a single microorganism which has been genetically engineered to produce ethanol as a primary fermentation product and to produce xylanase intracellularly. Specifically, the truncated xylanase gene (*xynZ*) from the thermophilic bacterium, *C. thermocellum*, was fused with the N-terminus of *lacZ* to eliminate secretory signals. This hybrid gene was expressed at high levels in the *E. coli* KO11 and *K. oxytoca* M5A1 (pLOI555) discussed above. Large amounts of xylanase accumulated as an intracellular product during ethanol production. Cells containing xylanase were harvested at the end of fermentation and added to a xylan solution at elevated temperature, thereby releasing xylanase for saccharification. After cooling, the hydrolysate was fermented to ethanol with the same organism thereby replenishing the supply of xylanase for a subsequent saccharification.

[0199] **Organism and growth conditions:** Bacterial strains and plasmids used in this example are listed in Table 12.

Table 12.

Bacteria strains and plasmids used		
Bacteria/plasmid	Characteristics	Source or reference
<i>E. coli</i> DH5 α	<i>lacZ</i> M15, <i>recA</i>	Bethesda Rsrch Lab
<i>E. coli</i> KO11	<i>frd</i> , <i>Cm</i> ^r , <i>lpe</i> ^{ts}	Ohta <i>et al.</i> ^c
<i>K. oxytoca</i> M5A1 (pLOI555)	<i>Cm</i> ^r , <i>pet</i> ^b	Example 15, <i>supra</i>
pCT1202	<i>Ap</i> ^r , <i>xynZ</i> ⁺	Grepinet <i>et al.</i> ^d
pLOI1001	<i>Ap</i> ^r , <i>xylB</i> ⁺	Utt <i>et al.</i> ^e
pLOI2000	<i>Ap</i> ^r , <i>xylB</i> ⁺	this example
pLOI2001	<i>Ap</i> ^r , <i>lacZ::xynZ</i> ⁺	this example

^b *pet* refers to the presence of *Z. mobilis pdc* and *adhB* genes on plasmid pLOI555.

^c Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57: 893-900.

^c Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57: 893-900.

^d Grepinet *et al.* [1988] *J. Bacteriol.* 170: 4582-4588.

^e Utt *et al.* [1991] *Appl. Environ. Microbiol.* 57: 1227-1234.

Table 12. (continued)

Bacteria strains and plasmids used		
Bacteria/plasmid	Characteristics	Source or reference
pLOI2002	Ap ^r , lacZ::xynZ ⁺	this example
pLOI2003	Ap ^r , lacZ::xynZ ⁺ , xylB ⁺	this example

[0200] Strains were grown in Luria broth supplemented with 50 g xylose/liter. Transformants of *E. coli* were selected on Luria agar plates containing 50 mg ampicillin/liter or 40 mg chloramphenicol/liter. Transformants of *K. oxytoca* M5A1 were selected on Luria agar containing 1000 mg ampicillin/liter or 40 mg chloramphenicol/liter.

[0201] Recombinant clones were screened for xylanase activity by using microtitre plates which contained 4-methylumbelliferyl-β-D-cellobiopyranoside (100 mg/liter) (Millet [1985] *FEMS Microbiol. Lett.* 29: 145-149). Similarly, expression of the *Butyrivibrio fibrisolvens* xylB gene encoding both xylosidase and arabinosidase activities was screened by incorporating 4-methylumbelliferyl-α-L-arabinofuranoside (20 mg/liter) into solid medium (Utt *et al.* [1991] *Appl. Environ. Microbiol.* 57: 1227-1234). Hydrolysis of these substrates by positive clones produced a fluorescent product (umbelliferone) which was readily detected under 340 nm ultra violet light.

[0202] **Genetic procedures and recombinant techniques:** Plasmid preparation, digestion with restriction enzymes, ligation, transformation, and gel electrophoresis were carried out using standard procedures. *E. coli* strain DH5α was used as the host for all plasmid constructions. Polymerase chain reactions were performed with the TempCycler Model 50 (Coy Laboratory Product Inc., Ann Arbor, MI) and a GeneAmp kit (Perkin Elmer Cetus, Norwalk, CT) containing Taq polymerase. Amplification reactions contained 2 mM each dNTP, 100 pmol each primer, 20 ng template, and 2.5 U Taq polymerase in 100 μl total volumes. Products were isolated after 30 cycles of amplification (1 minute at 94°C, 2 minutes at 47°C and 1 minute at 72°C; final extension for 3 minutes at 72°C).

[0203] **Determination of enzymatic activities:** Xylanase, xylopyranosidase, and arabinofuranosidase activities were measured in recombinants which had been grown overnight at 30°C. Cells were harvested by centrifugation (7000 x g, 10 minutes) and washed twice with phosphocitrate buffer (50 mM potassium phosphate and 12.5 mM citric acid, pH 6.3) for the determination of xylanase activity or with phosphate buffer (5 mM sodium phosphate buffer containing 10 mM 2-mercaptoethanol, pH 6.8) for the measurement of xylosidase and arabinosidase activities. Cells were disrupted by two passages through a French pressure cell at 20,000 psi. The resulting lysate was centrifuged (13,000 x g for 30 minutes, 4°C) to remove cell debris.

[0204] Xylosidase and arabinofuranosidase activities were assayed as described previously (Utt, *supra.*). Xylanase was measured by the same procedure but using p-nitro-β-D-cellobioside as the substrate. Xylanase activity was also estimated by measuring the release of reducing sugars (Bergmeyer (ed) [1983] *Methods of enzymatic analysis*, Vol. II, 3rd edition, p 151-2, Verlag Chemie, Weinheim, Germany) from the hydrolysis of birchwood xylan (Sigma Chemical Company, St. Louis, MO). All activities are reported as millimicromoles of product liberated per minute. Protein concentration was determined by the method of Bradford (Bradford [1976] *Anal. Biochem.* 72: 248-254).

[0205] **Separation of xylooligosaccharides by thin layer chromatography:** A mixture of xylooligosaccharides was prepared by the partial hydrolysis of 0.5% birchwood xylan with trifluoroacetic acid (Dorner [1988] *Meth. Enzymol.* 160: 176-180). The solution was concentrated 10-fold under vacuum at room temperature and 1 μliter samples used as a standard. Xylooligosaccharides were separated at room temperature by single development on unactivated Whatman silica gel 150A plates (Whatman Inc., Clifton, New Jersey) using a solvent composed of acetone, ethylacetate, and acetic acid (2:1:1, respectively). After drying, oligosaccharides were visualized with naphthylethylenediamine reagent as described by Bounias (Bounias [1980] *Anal. Biochem.* 106:291-295).

[0206] **Fermentation experiments:** Inocula for fermentations were prepared from freshly isolated colonies by overnight growth in unshaken flasks (30°C). Fermentations were inoculated to an initial O.D. at 550 nm of either 0.1 or 0.3 and incubated at 30°C in stirred, pH-stats (350 ml working volume, pH 6.0). (Beall *et al.* [1991] *Biotechnol. Bioengin.* 38: 296-303.; Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57: 893-900; and Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57:2810-2815.) Chloramphenicol or chloramphenicol and ampicillin were included in fermentation broths.

[0207] Xylan was fermented by a two stage process in which xylan degradation was carried out at elevated temperature followed by fermentation at 30°C. For the degradation of xylan, cells from 350-ml fermentations were harvested by centrifugation (7,000 x g, 10 minutes) after 48 hours, resuspended in an equal volume of fresh Luria broth containing 4% xylan (pH 6.0), and incubated at 60°C for 65 hours. After cooling to 30°C, hydrolysates were inoculated with recombinant organisms to provide an initial O.D. of 0.3 at 550 nm.

[0208] Samples were removed at various times to monitor xylooligosaccharides (thin layer chromatography), cell growth (O.D. at 550 nm), and ethanol. Ethanol was measured by gas liquid chromatography (Dombek *et al.* [1985] *Appl. Environ. Microbiol.* 51:197-200).

[0209] **Construction of recombinant plasmids for the hydrolysis of xylan:** At least two activities are required for

the hydrolysis of unsubstituted xylan, i.e., an endolytic xylanase and xylosidase. Three plasmids were constructed for us in xylan fermentations in which the *xynZ* (xylanase) gene from the thermophile, *Clostridium thermocellum* (Grepinet *et al.* [1988] *J. Bacteriol.* 170: 4582-4588), and the *xyiB* (xylosidase and arabinosidase) gene from *Butyrivibrio fibrisol-*

vens (Utt, *supra.*) were expressed singly and in combination as an operon (Figs. 7A, 7B and 7C).
[0210] Figures 7A, 7B, and 7C illustrate the construction of recombinant plasmids containing *xynZ* and *xyiB*. Coding regions are boxed. *C. thermocellum* and *B. fibrisolvens* DNA are represented by thin lines. The thick line denotes vector DNA. Sites of blunt-ended ligation are marked by an "X." An "F" labels the coding region for the in-frame *lacZ::xynZ* fusion.

[0211] The *xyiB* gene was subcloned into the *XbaI* to *PstI* region of the polylinker in pUC18 by inserting two fragments from pLOI1001, a 0.3 kilobase-pair (kbp) *XbaI* to *PstI* fragment containing the amino terminus with ribosomal-binding site and a 2.4 kbp *PstI* fragment containing the remainder of the coding region and translational terminator. The resulting plasmid, pLOI2000, expressed the *xyiB* gene from the *lac* promoter.

[0212] Previous studies by Grepinet *et al.*, *supra.*, have shown that xylanase expression was elevated in *lacZ* fusions in which a large segment encoding the secretion signal sequence and amino terminus of the processed xylanase had been deleted. A very similar *lacZ::xynZ* fusion was constructed by blunt-ended ligation of the Klenow-treated 2.4 kbp *StyI* fragment from pCT1202 containing the amino-truncated *xynZ* gene into the Klenow-treated *PstI* site of pUC18. Both reading frames were aligned in the resulting plasmid, pLOI2001.

[0213] It was necessary to remove the putative transcriptional terminator 30 bp downstream from the coding region of *lacZ::xynZ* and add a new *SstI* site to the 3'-end of the fusion gene prior to constructing a plasmid from which genes encoding both xylanase and xylosidase were expressed. These modifications were made by using polymerase chain reaction. Plasmid pLOI2001 was used as the template with 5'-GAATTCGAGCTCGGTAC-3' as a primer for the 5'-end and 5'-GGGAGCTCCGGCATCATTATCTG-3' as a primer for the 3'-end (includes a new *SstI* site). After *SstI* digestion, this fragment was inserted into the *SstI* polylinker sites of pUC18 and pLOI2000 to construct pLOI2001 and pLOI2003, respectively.

[0214] **Expression of enzymes involved in xylan degradation:** The expression of xylanase, xylosidase, and arabinosidase activities were initially examined in stationary phase cells of the host organism used for constructions, strain DH5a (Table 13). Xylanase (*xynZ*) activity was reduced by 60% in clones harboring pLOI2003 which contains *xyiB* downstream as compared to pLOI2001 which contains *xynZ* alone. However, xylosidase and arabinosidase (*xyiB*) activities were similar in clones harboring pLOI2000 containing *xyiB* alone and pLOI2003 containing an upstream *xynZ* gene.

Table 13.

Specific activities of recombinant enzymes for xylan degradation				
Strain and plasmid	Specific activity (mU/mg)			
	xylosidase	arabinosidase	xylanase ^a	xylanase ^b
E. coli DH5α	0	0	0	0
pLOI2000	1.2	2.2	0	0
pLOI2001	0	0	1.4	124
pLOI2003	1.5	2.5	0.5	48
E. coli KO11	0	0	0	0
pLOI2001	0	0	0.4	38
pLOI2003	1.3	2.4	0.3	25
pLOI2003 ^c	1.1	1.9	0.3	47
pLOI2003 ^{c,d}	nd	nd	0.8	93
K. oxytoca M5A1 (pLOI555)	53	0	0	0
& pLOI2001	49	0	0.4	39
pLOI2003	56	2.6	0.2	24
pLOI2001 ^c	30	0	0.7	0
pLOI2001 ^{c,d}	nd	nd	1.8	144

^a Xylanase activities were measured using p-nitrophenyl-β-D-cellobioside as a substrate.

^b Xylanase activities were measured using 0.5% birchwood xylan as a substrate.

^c Cells were harvested from pH-stats (8% xylose in Luria broth, pH 6.0) at the end of fermentation.

^d Xylanase activities were determined at 60°C.

Table 13. (continued)

Specific activities of recombinant enzymes for xylan degradation				
Strain and plasmid	Specific activity (mU/mg)			
	xylosidase	arabinosidase	xylanase ^a	xylanase ^b
pLOI2003 ^c	38	2.9	0.4	58
pLOI2003 ^{c,d}	nd	nd	0.8	144

Unless indicated otherwise: cells were grown in shake flasks containing Luria broth with 5% xylose and harvested after 20 hours (stationary phase); xylosidase and arabinosidase activities were determined at 30°C; and xylanase activities were measured at 45°C.

^a Xylanase activities were measured using p-nitrophenyl-β-D-cellobioside as a substrate.

^b Xylanase activities were measured using 0.5% birchwood xylan as a substrate.

^c Cells were harvested from pH-stats (8% xylose in Luria broth, pH 6.0) at the end of fermentation.

^d Xylanase activities were determined at 60°C.

[0215] The plasmids pLOI2001 and pLOI2003 were transformed into the ethanologenic strain of *E. coli* KO11, in which the genes for ethanol production are integrated into the chromosome. Expression was compared in stationary phase cells from shake flasks and in cells from a pH-stat at the end of fermentation (Table 13). Xylosidase, arabinosidase, and xylanase activities were equivalent to those observed with DH5α. Again, xylanase activity was reduced in pLOI2003 in which xylB was present downstream.

[0216] Expression was also examined in *K. oxytoca* strain M5A1 harboring pLOI555 containing genes from the *Z. mobilis* ethanol pathway. Although the type of replicon present in pLOI555 is unknown, it appeared quite stable in the presence of a second plasmid constructed from pUC18. Expression of arabinosidase and xylosidase activities were roughly equivalent to that observed in *E. coli*. An abundant native xylosidase was discovered in M5A1. Xylosidase activities were somewhat higher in cells from shake flasks as compared to cells from pH-stats while xylanase activities followed the opposite trend.

[0217] Under the assay conditions used, arabinosidase activity was 1.5 to 1.7 times higher than xylosidase activity confirming previous reports (Utt *et al.*, *supra*). Xylanase activity measured at 47°C was approximately half that measured 60°C, the optimal temperature for the native enzyme (Grepinet *et al.*, *supra*). Based on the data presented, specific activities computed from reducing sugar assays for the xylanase fusion are approximately 100-fold higher than estimates based on the hydrolysis of p-nitrophenyl-β-D-cellobioside indicating a strong preference for native substrate.

[0218] **Hydrolysis of xylan at elevated temperature by recombinant clones:** Strain KO11(pLOI2003) was grown in a pH-stat containing 8% xylose and tested as a source of enzyme for xylan hydrolysis. Cells were resuspended in an equal volume of fresh Luria broth containing 4% xylan and incubated at 45°C, the maximal temperature at which *B. fibrisolvens* xylosidase is stable, and 60°C and the optimal temperature for *C. thermocellum* xylanase. Although xylanase and xylosidase were produced as intracellular products, initial experiments indicated that these enzymes were readily released into the medium by incubation at 45° and 60°C. Samples were removed at various times and digestion products analyzed by thin layer chromatography (Figs. 8A and 8B).

[0219] Figures 8A and 8B show thin layer chromatographic analysis of xylan hydrolysis using *E. coli* strain KO11 (pLOI2003) incubated at 45°C (Fig. 8A) or 60°C (Fig. 8B) in Luria broth (pH 6.0) containing 4% birchwood xylan. A sample size of 1 μliter was applied to each lane. Incubation times are given in hours beneath each lane. An acid hydrolystate of xylan was used as a standard in the first lane (S).

[0220] Xylose, xylobiose, xylotriose, and xylotetrose were clearly resolved with xylobiose being the dominant product. Monomeric xylose accumulated slowly. A comparison of the extent of digestion after 24 and 48 hours clearly shows that less hydrolysis occurred at 45°C than at 60°C despite the instability of xylosidase at 60°C.

[0221] Although xylan digestion was incomplete even after 65 hours at 60°C, xylanase remained quite active and was readily detected with 4-methyl-umbelliferyl-β-D-cellobioside as a substrate. Xylosidase was rapidly inactivated at this temperature. No further change in thin layer profiles of xylan hydrolysate were observed after the addition of cell lysates containing both enzymes and incubation at 60°C for another 24 hours. Thus, the xylooligosaccharide products appear to be near limit digestion products of Sigma birchwood xylan. Oxidation products or substitutions may block complete digestion of this commercial preparation.

[0222] **Utilization of xylan hydrolysis products by recombinant strains of *E. coli* and *K. oxytoca*:** Small scale experiments were conducted to evaluate the extent to which xylooligosaccharides could be metabolized by *E. coli* KO11(pLOI2003) and *K. oxytoca* strain M5A1(pLOI555). These strains were inoculated into 1 ml of xylan hydrolysate (60°C for 65 hours; supernatant after removal of cell debris by centrifugation) and incubated for 48 hours at 30°C without agitation. Samples were removed and analyzed by thin layer chromatography (Figs. 9A and 9B).

[0223] Figures 9A and 9B show the utilization of xylooligosaccharides for growth by recombinant *E. coli* KO11 (Figure 9A) and *K. oxytoca* M5A1 (Figure 9B). Xylan (4%) was hydrolyzed for 65 hours by incubation with *E. coli* strain KO11 (pLOI2003) in Luria broth (pH 6.0), centrifuged, and sterilized by filtration. Samples were inoculated and incubated at 30°C to allow growth. Samples were removed at 24 hours intervals as indicated beneath each lane and analyzed by thin layer chromatography. A mixture of xylooligosaccharides was used as a standard in the first lane (S).

[0224] Only xylose was metabolized by the recombinant *E. coli* despite the presence of active *B. fibrisolvens* xylosidase. Xylose, xylobiose and xylotriose were all completely consumed by the ethanologenic *K. oxytoca*. Based upon these results, derivatives of *K. oxytoca* were selected as being superior for further studies of xylan conversion to ethanol.

[0225] **Fermentation of xylose and xylan by derivatives of *K. oxytoca* strain M5A1:** Fermentation of xylan was carried out as a two-stage process in which cells harvested from a previous fermentation were resuspended in Luria broth containing 4% xylan for saccharification (60°C for 65 hours). Saccharified xylan was inoculated and fermented at 30°C (pH 6.0). Parallel experiments were conducted in which the cell debris had been removed by centrifugation before inoculation. Fermentations were also conducted with 4.47% xylose (equivalent to the xylose content of 4% xylan) for comparison (Figs. 10A and 10B). The data from those fermentations is provided below in Table 14.

[0226] Figures 10A and 10B show the conversion of xylose and xylan hydrolysate to ethanol by recombinant *K. oxytoca* M5A1. Xylan was hydrolyzed for 65 hours at 60°C using cells from a previous fermentation as the source of enzymes. Figure 10A shows the growth of the cell mass, and Figure 10B shows the ethanol production. Symbols: ▲, fermentation of 4.47% xylose by M5A1(pLOI555); ○, fermentation of 4.47% xylose by M5A1(pLOI555 and pLOI2001); ■, fermentation of 4% xylan hydrolysate containing cell debris from saccharification by M5A1(pLOI555 and pLOI2001); □, fermentation of 4% xylan hydrolysate after centrifugation to remove cell debris by M5A1(pLOI555 and pLOI2001).

Table 14.

Ethanol production from xylose and xylan by recombinant strains of <i>K. oxytoca</i> M5A1(pLOI555) ^a							
Substrate and second plasmid	Base (mM/g of sugar) ^b	Time ^c (h)	Ethanol Yield		Theoretical yield (%) ^d	VP (g/liter/h) ^e	Cell Yield (g/g of sugar)
			g/liter	g/g of substrate			
<u>Xylose (4.47%)</u>							
pLOI555 alone	1.2	36	22.6	0.51	99	1.29	0.07
with pLOI2001	1.5	48	21.7	0.49	95	1.02	0.08
with pLOI2003	1.3	36	20.6	0.47	91	0.83	0.07
<u>Xylan (4.0%)</u>							
with pLOI2001 ^f	0.7	24	7.7	0.19	34	0.37	0.04
with pLOI2003 ^f	1.3	36	7.7	0.19	34	0.31	0.04
with pLOI2001 ^g	1.3	60	7.8	0.20	34	0.28	nd
with pLOI2003 ^g	1.1	60	7.9	0.20	35	0.30	nd

^a Calculations are based on total substrate added.

^b Amount of base consumed to maintain a pH of 6.0 during fermentation.

^c Time required to reach maximal ethanol concentration.

^d Theoretical yield on a weight basis is 0.51 for xylose and 0.56 for xylan.

^e VP, maximum volumetric productivity during batch fermentation.

^f Hydrolysate after debris from saccharification was removed by centrifugation.

^g Hydrolysate containing debris from saccharification.

[0227] It is seen that M5A1(pLOI555) produced ethanol very efficiently. Xylose fermentation was essentially completed after 24 hours with 93% of the maximal theoretical yield. Both growth and ethanol production from xylose were reduced by the simultaneous presence of a second plasmid, either pLOI2001 or pLOI2003. Required fermentation times with

pLOI2001 increased to approximately 36 hours with 91% of theoretical yield. This decrease in ethanol productivity from monomeric sugar reflects the additional burden imposed by the recombinant enzymes.

[0228] Xylan hydrolysis (60°C for 65 hours) was carried out with M5A1(pLOI555) containing either the hybrid xylanase gene (pLOI2001) or the xylosidase and hybrid xylanase gene (pLOI2003). Thin layer profiles of these hydrolysates appeared identical to those of KO11 (pLOI2003), shown in Figs. 9A and 9B. No differences were found in the extent of xylan hydrolysis between the two M5A1 derivatives.

[0229] Growth and ethanol production were measured in these hydrolysates after inoculation with the respective organism used in saccharification (Figs. 10A and 10B). Growth in clarified hydrolysate was only half that observed with equivalent levels of xylose monomer. Fermentations were essentially complete after 24 hours and appeared slightly faster in clarified hydrolysates. Ethanol yields from xylan were approximately 1/3 of maximal theoretical values, 7.7 to 7.9 g/liter.

[0230] Xylooligosaccharides in fermentation broth were monitored by thin layer chromatography. The profiles at the end of fermentation were identical to that shown in Fig. 9B (48 hours). Xylose, xylobiose, and xylotriose were completely metabolized with xylotetrose and longer oligomers remaining.

[0231] **Discussion:** It is thus seen that highly engineered strains of *K. oxytoca* M5A1 can be used both as a source of enzyme for polymer hydrolysis and for ethanol production from xylan. The presence of a native, intracellular xylosidase and the ability to transport and metabolize xylooligosaccharides have not been previously reported for this organism. *E. coli* isolates from nature have been previously described which utilize a phosphotransferase system for cellobiose and an intracellular phosphocellobiase (Hall *et al.* [1987] *J. Bacteriol.* 169: 2713-2717; Kricker *et al.* [1987] *Genetics* 115: 419-429). Analogous systems may function in *K. oxytoca* M5A1 for xylobiose and xylotriose metabolism. The presence of transport systems and enzymes for the metabolism of xylose monomers, dimers and trimers in *K. oxytoca* M5A1 offers considerable advantage over *E. coli*-based or *S. cerevisiae*-based systems for further development of biomass to ethanol processes.

[0232] Many of the previous problems in the genetic engineering of single organisms for polymer degradation and ethanol production are eliminated by using a two-staged process with thermostable enzymes being produced as intracellular products during fermentation. Synthesis of proteins destined to be secreted at high levels often adversely affect normal cellular processes. This problem should be minimized by using N-truncated enzymes in which the secretory signal has been deleted for intracellular expression. Since hydrolysis time is a limiting factor in conversion, high levels of hydrolytic enzymes would be most advantageous at the very beginning of the conversion process to maximize early release of sugars for rapid fermentation. In secretory-based processes, enzyme levels are initially low, accumulating to reach maximal levels near the end of fermentation. However, by using the cells from previous fermentations as a source of enzymes one achieves near maximal levels initially. Although the use of thermostable enzymes for hydrolysis offers additional process advantages for minimizing contamination and may provide higher rates of hydrolysis, a primary advantage is the simplicity of release of intracellular enzymes from harvested cells.

[0233] Ethanol tolerance is frequently a problem in native organisms which degrade cellulose and xylan. Such organisms typically produce less than 20 grams of ethanol per liter. (See Tailleux *et al.* [1989] *Appl. Environ. Microbiol.* 55: 203-206.) Although high level ethanol production from xylan thus far has not been achieved, M5A1(pLOI555) is capable of producing at least 48 g ethanol/liter from 100 g/liter xylose (approximately 95% of maximal theoretical yield).

[0234] The overall yield of ethanol from birchwood xylan was lower than anticipated and is believed to be primarily due to incomplete xylan hydrolysis. Undegraded xylooligosaccharides remained after the 65 hours saccharification period. Although the level of xylanase produced by our constructs was below that produced by optimal xylanase fusion constructs reported by Grepinet *et al.*, *supra*, enzyme level *per se* did not appear limiting. Xylan (40 g/liter; approximately 0.3 moles of anhydroxylose) should be completely degraded into xylobiose equivalents after 24 hours by hydrolysis at a rate of 105 μ moles/min. Cell yields from fermentation were approximately 4.5 g cell protein/liter with a xylanase activity of 144 μ moles/min per g cell protein providing a total activity of 648 μ moles/min. Xylooligosaccharides remained, however, even after 65 hours despite the persistence of active xylanase. These levels of xylooligosaccharides were unchanged by the addition of fresh enzyme and further incubation for 24 hours. Competitive inhibition of hydrolysis by xylooligosaccharides may contribute to incomplete digestion and was tested, in part, by hydrolysis at 45°C, a temperature at which xylanase remained active as did the native enzymes in M5A1 which metabolized xylose, xylobiose, and xylotriose. The oligosaccharide profile under these conditions was identical to that observed after fermentation by M5A1 strains with equivalent levels of the xylotetrose and longer oligosaccharides remaining undigested (data not shown). These oligosaccharides appear to be limit digestion products although the nature of the substituents limiting digestion remains unknown. According to manufacturer, the birchwood xylan is 99% xylose. While not wishing to be bound to any particular theory, the present inventors speculate that this product may contain oxidized residues generated during storage or substitutions which have survived base extraction and purification (Chesson *et al.* [1983] *J. Sci. Food Agric.* 34: 1330-1340).

[0235] Fermentative ethanol production in the U.S. is approximately 1 billion gallons (3.8 billion liters) per year (Lynd *et al. Science* 251:1318-1323). Microorganisms such as the genetically engineered *K. oxytoca*, *E. coli*, *S. cerevisiae*,

or *Z. mobilis* could be used to produce a variety of enzymes as coproducts with ethanol. At a minimal cell yield of 2 grams of protein/liter of beer, the diversion of 5% of cell protein would produce at least 3,800,000 kg of enzymes as co-products. These enzymes need not be limited to depolymerization of substrates for fermentation but could also include other enzymes for large markets such as the detergent industry, the food industry, the pulping of wood, and the development of new biocatalyst-based industries for novel chemicals. Providing large markets develop, the value of such enzymes could well exceed current values of ethanol itself.

Example 18 - Ethanol Production from Cellobiose, Amorphous Cellulose, and Crystalline Cellulose by Recombinant *Klebsiella oxytoca* containing Chromosomally Integrated *Z. mobilis* Genes for Ethanol Production and Plasmids Expressing Thermostable Cellulase genes from *Clostridium thermocellum*

[0236] Summary of Example: In this experiment, the *Z. mobilis* genes for ethanol production have been integrated into the chromosome of *Klebsiella oxytoca* strain M5A1. The best of these constructs, strain P2, produced ethanol efficiently from cellobiose in addition to monomeric sugars. Utilization of cellobiose and cellotriose by this strain eliminated the requirement for external β -glucosidase and reduced the amount of commercial cellulase needed to ferment SOLKA FLOC SW40 (primarily crystalline cellulose). The addition of plasmids encoding endoglucanases from *Clostridium thermocellum* resulted in the intracellular accumulation of thermostable enzymes as co-products with ethanol during fermentation. The best of these, strain P2(pCT603T) containing *celD*, was used to hydrolyze amorphous cellulose to cellobiose and produce ethanol in a two stage process. Strain P2(pCT603T) was also tested in combination with commercial cellulases. Pretreatment of SOLKA FLOC SW40 at 60°C with endoglucanase D reduced the amount of commercial cellulase required to ferment SOLKA FLOC by up to 80%. The stimulatory effect of the endoglucanase D pretreatment may result from the hydrolysis of amorphous regions, exposing additional sites for attack by fungal cellulases. Since endoglucanase D functions as part of a complex in *C. thermocellum*, it is possible that this enzyme may complex with fungal enzymes or bind cellulose to produce a more open structure for hydrolysis.

[0237] Example 17 demonstrated the advantage of accumulating the *Clostridium thermocellum* xylanase intracellularly in *K. oxytoca* M5A1 as a co-product with ethanol. In this example, the present inventors have integrated the genes for ethanol production into *K. oxytoca* strain M5A1 and demonstrated that the resulting organism can efficiently convert cellobiose to ethanol. This organism appears capable of transporting and metabolizing both cellobiose and cellotriose, eliminating the need for heterologous β -glucosidase. Plasmids added to this integrated construct allowed the production of thermostable endoglucanases as co-products with ethanol, reducing the requirement for commercial cellulases during the fermentation of cellulose.

[0238] Organisms and growth conditions: Table 15 lists the new organisms and plasmids used in this study. Strains harboring ethanologenic genes were maintained on Luria agar containing 2% glucose (3); other strains were maintained on Luria agar lacking added sugar. Unless noted otherwise, chloramphenicol (20 μ g/ml), tetracycline (6 μ g/ml), and ampicillin (50 μ g/ml) were used for selection. Typically, *E. coli* strains were grown at 37°C and *K. oxytoca* strains grown at 30°C. Expression of the ethanologenic operon from *Z. mobilis* was monitored by the rate of color formation on aldehyde indicator plate (Conway et al. [1987] *J. Bacteriol.* 169:2591-2597). Colonies were screened for cellulase activity by incubating for 1 to 2 hours at 55°C on Luria agar containing carboxymethyl cellulose followed by staining with Congo Red (Wood et al. [1988] *Meth. Enzymol.* 160:59-74).

TABLE 15.

Bacteria strains and plasmids used in this study		
Bacteria strains/plasmid	Characteristics	Source or reference
strains		
<i>K. oxytoca</i> M5A1(pLOI555)	Cm ^r , <i>pet</i> ^b	Example 15
S1	Cm ^r , <i>lpet</i> ^a	this example
S2	Cm ^r , <i>lpet</i> ^a	this example
S3	Cm ^r , <i>lpet</i> ^a	this example
P1	Cm ^r , <i>lpet</i> ^a	this example
P2	Cm ^r , <i>lpet</i> ^a	this example
B1	Cm ^r , <i>lpet</i> ^a	this example
plasmids		
pCOS2EMBL	Tc ^r	Poustka <i>et al.</i> ^c
pLOI510	Cm ^r <i>pet</i> ^b	Ohta <i>et al.</i> ^d
pCT105	Ap ^r <i>celA</i> ⁺	Comet <i>et al.</i> ^e
pCT207	Ap ^r <i>celB</i> ⁺	Jeffries ^f
pCT301	Ap ^r Tc ^r <i>celC</i> ⁺	Petre <i>et al.</i> ^g
pCT603	Ap ^r Tc ^r <i>celD</i> ⁺	Joliff <i>et al.</i> ^h
pCT105T	Ap ^r Tc ^r <i>celA</i> ⁺	this example
pCT207T	Ap ^r Tc ^r <i>celB</i> ⁺	this example
pCT603T	Ap ^r Tc ^r <i>celD</i> ⁺	this example

^a *lpet* refers to the integration of *Z. mobilis* *pd*c and *adhB* genes into the chromosome.

^b *pet* refers to presence of *Z. mobilis* *pd*c and *adhB* genes on plasmid pLOI555.

^c Poustka *et al.* [1984] *Proc. Natl. Acad. Sci. USA* 81:4129-4133.

^d Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57:2810-2815.

^e Comet *et al.* [1983] *Bio/Technology* 1:589-594.

^f Jeffries, in J.F. Kennedy *et al.* (ed.), *Wood and Cellulose: Industrial utilization, biotechnology, structure and properties*. John Wiley & Sons, New York (1988).

^g Petre *et al.* [1986] *Biochimie* 68:687-695.

^h Joliff *et al.* [1986] *Bio/Technology* 4:896-900.

[0239] Genetic methods and plasmid construction: Standard genetic procedures were used. Integration of *pet* genes into strain M5A1 was obtained by transforming cells with circularized DNA (lacking all replication functions) containing three essential elements: 1) *E. coli* *pfl* gene or M5A1 DNA; 2) the *Z. mobilis* *pet* genes; 3) the *cat* gene for selection. Recombinants were initially selected using 20 µg chloramphenicol/ml and expressed low levels of *Z. mobilis* enzymes. As with *E. coli* (Ohta *et al.* [1991] *Appl. Environ. Microbiol.* 57: 893-900), expression was boosted by direct selection of mutants with resistance to 600 µg chloramphenicol/ml. A single clone expressing high level resistance was retained for each independent integration event.

[0240] Tetracycline resistance was added to plasmids containing *celA* (pCT105), *celB* (pCT207), and *celD* (pCT603). Plasmids pCT105 and pCT207 were digested with *Bam*HI followed by treatment with the Klenow fragment of *E. coli* DNA polymerase to create blunt ends. Plasmid pBR322 was digested with *Eco*RI and similarly treated to produce blunt ends. All three plasmids were then digested with *Sal*I. The resulting small fragment from pBR322 containing the 5'-end of *tet* was ligated to the larger fragments from pCT105 and pCT207 to reform a functional *tet* gene (previously inactivated by insertion of *C. thermocellum* DNA), denoted pCT105T and pCT207T, respectively. A 2.5 kilobase pair (kbp) *Eco*RI fragment from pCOS2EMBL (Poustka *et al.*, *supra*) containing the complete gene was inserted into the unique *Eco*RI in pCT603 to produce pCT603T. No modification of pCT301 was needed since this plasmid contained a functional *tet* gene in addition to *celC*. All cellulase-containing plasmids were transformed into recombinant *K. oxytoca* with selection for tetracycline resistance.

[0241] Cellulase activity: Endoglucanase activity was determined in cells plus broth and in broth alone at 60°C with p-nitrophenyl-β-D-cellobioside as a substrate (Petre *et al.*, *supra*). Endoglucanase D activity was also estimated as the release of reducing sugars from amorphous cellulose. Amorphous cellulose (acid swollen and base swollen) were prepared from SOLKA FLOC SW40 as previously described for Avicel (Wood [1988] *Meth. Enzymol.* 160:19-25). Reducing sugars were measured by using the Nelson-Somogyi method (Bergmeyer *et al.* [1983] pp 151-152, in Bergmeyer H.U. (ed), *Methods of enzymatic analysis*, vol. II, 3rd edition, Verlag Chemie, Weinheim, Germany). Initial concentra-

tions of amorphous cellulose were estimated as total carbohydrate by the phenolsulfuric acid method (Wood [1988] *Meth. Enzymol.* 160:87-116).

[0242] Endoglucanase activities were also compared among ethanologenic *cel⁺* recombinants by overlaying 18 mm x 150 mm culture tubes containing approximately 15 ml of Luria broth solidified with 10% low viscosity carboxymethyl cellulose (Sigma Chemical Company, St. Louis, Mo.) with 2 ml of stationary phase culture. After 48 hours of incubation at 55°C, the extent of liquefaction was determined by inversion.

[0243] Products of cellulose digestion were analyzed by thin layer chromatography as described previously for xylosides and xylan digestion (Domer *et al.* [1988] *Meth. Enzymol.* 160:355-362). Glucose, cellobiose, and cellotriose were separated from oligomers. Cellobiose and glucose served as standards.

[0244] **Fermentation experiments:** Fermentations were carried out in 500-ml Fleakers which served as pH stats (350 ml working volume) essentially as described previously (Beall *et al.*, *supra*). Luria broth containing either 10% glucose or 10% cellobiose was tested at 30°C, pH 6.0, 100 rpm. Inocula for fermentations were grown at 30°C overnight in unshaken, 250-ml flasks containing 50 ml of Luria broth (4% glucose). After mixing, cell densities were measured at 550 nm and used to calculate the volume required to provide an initial density of 0.32 mg cell dry weight/ml (O.D. 550 nm of 1.0). Cells were harvested from this volume by centrifugation and resuspended in a portion of the broth from each respective pH-stat to start fermentation.

[0245] Sugars were sterilized separately by filtration. Cellulose fermentations contained 50 g/liter SOLKA FLOC SW40 (James River Corporation, Saddle Brook, NJ) and were carried out at 35°C. Cellulose was sterilized by autoclaving as a dry powder. For investigations of cellulose fermentation using commercial enzymes, CYTOLASE or MULTIFECT, was added at the time of inoculation.

[0246] Samples were removed for the determination of cell mass (O.D. 550 nm) and ethanol (gas liquid chromatography; Beall *et al.*, *supra*). Ethanol concentrations are expressed as g/liter. Ethanol yields were corrected for dilution by the addition of base during fermentation and computed on the basis of total sugar or cellulose initially present. No corrections were made for residual carbohydrates. Maximum theoretical yields from glycolysis and fermentation are 0.51 g ethanol/g glucose, 0.536 g ethanol/g cellobiose, and 0.56 g ethanol/g cellulose. Except as indicated, results presented are an average of 2 or more fermentations.

[0247] **Commercial cellulases:** Two commercial cellulases were investigated, CYTOLASE M104 and MULTIFECT CL (Genencor International, Rolling Meadows, IL). Both were supplied as liquids, presumably broths from mutant *Trichoderma longibrachiatum* with proprietary amendments. Both are described as containing a mixture of pectinases, cellulases, and hemicellulases. These enzyme preparations are currently optimized for food processing.

[0248] **Chromosomal integration of *Z. mobilis* *pet* genes into M5A1:** Three approaches were used to integrate *pet* genes into the chromosome of M5A1 (Orsdov [1984] p. 461-465, in N.R. Kreig and J.G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 1. The Williams and Wilkins Co., Baltimore). Since *K. oxytoca* and *E. coli* are closely related, the *E. coli* *pfl* gene was used as a potential source of homologous DNA to promote recombination analogous to the method used for *E. coli* (Ohta *et al.* *Appl. Environ. Microbiol.* 57:893-900). *SacI* fragments of 8.6 kbp were purified from pLOI510 which contained *pet* genes and *cat* within the *pfl* gene of *E. coli*. This fragment (lacking genes involved in replication) was circularized by ligation and transformed into M5A1 to allow direct selection for integration. A shorter 5 kbp *PstI* fragment containing only a small amount of flanking *E. coli* *pfl* was used in a similar manner. In the third approach, homologous *K. oxytoca* M5A1 DNA (2 kbp random *Sau3A* fragments) was ligated to a 4.6 kbp *Bam*HI fragment containing only the *pet* genes with *cat* (no *E. coli* DNA) and used for transformation. Integrated strains were recovered from all three approaches by selection on Luria broth plates containing 2% glucose and 20 µg Cm/ml. Three integrated clones were recovered with the *SacI* fragment, two with the *PstI* fragment and one with the *Bam*HI fragment.

[0249] After overnight growth in liquid culture, 0.1 ml of stationary phase cells was spread on plates containing 600 µg Cm/ml and 2% glucose to select for high level expression. Single large colonies were retained from each independent integration event and named according to the restriction site used for construction, i.e., S1, S2, S3, P1, P2, and B1. These clones were tested for the presence of plasmids by transformation of miniscreen DNA and for expression of *pet* genes on aldehyde indicator plates (Table 16). Putative integrated strains found to contain plasmid-borne *cat* genes were discarded after digesting miniscreen DNA to confirm the presence of pLOI510 (presumably a low level contaminant of gel-purified fragments). The parent organism and M5A1(pLOI555), an excellent ethanol producer, were included as negative and positive controls, respectively. Two clones expressed the ethanol genes at levels nearly equivalent to that of that of M5A1(pLOI555), strains B1 and P2.

TABLE 16.

Fermentation of 10% glucose (48 hours) by strains of M5A1 containing integrated <i>pet</i> genes.					
Strain ^a	Base (mM/liter)	Cell Mass ^b (g/liter)	Ethanol (g/liter)	Aldehyde ^c Plates	Plasmid ^d
M5A1	108	3.2	15	-	-
M5A1 (pLOI555)	91	5.1	45	++++	+
S1	120	2.9	37	+	-
S2	171	3.2	44	++	-
S3	97	2.4	37	++	-
P1	120	2.0	38	++	-
P2	91	3.7	44	+++	-
B1	63	4.0	47	++++	-

^a Values reported are from a single experiment although many were reproduced and averaged in subsequent investigations.

^b Cell mass was calculated from the maximal O.D. at 550 nm (approx. 0.32 g dry weight/liter per O.D. unit).

^c Comparative rate of color development on aldehyde indicator plates at 30°C was used as a relative measure of expression from the *pet* operon.

^d Small scale preparations of DNA were examined on agarose submarine gels and tested for their ability to transfer antibiotic resistance to *E. coli* DH5α during transformation.

[0250] Comparison of glucose fermentation by integrated strains to M5A1(pLOI555): After 48 hours, high levels of ethanol equivalent to M5A1(pLOI555) were produced by 4 strains containing *pet* integrations (Table 16). The two which produced the lowest levels of acidic co-products also grew to the highest densities. These strains, P2 and B1, were selected for further investigation.

[0251] Fermentation of glucose and cellobiose by strains P2 and B1. Strains P2 and B1 were investigated for their ability to ferment 10% glucose and 10% cellobiose (Table 17). During glucose fermentation (Fig. 11A), both integrated strains produced 3 times more ethanol than the parent organism, M5A1, but were slightly inferior to M5A1 (pLOI555) (plasmid-borne *pet* genes) in yield, and in volumetric productivity. Unexpectedly, integration and high level expression of *pet* genes in strain B1 was accompanied by a loss of ability to ferment cellobiose. Strain P2, however, grew well and produced ethanol from cellobiose (Fig. 11B) with 96% of the maximum theoretical yield. Strain P2 was deposited as ATCC 55307 on March 11, 1992.

[0252] Figures 11A and 11B thus illustrate the ethanol production by recombinant strains of *K. oxytoca* M5A1. Figure 11A illustrates the production from glucose (100 g/liter). Symbols: ●, strain M5A1(pLOI555); ▲, strain P2 containing integrated *pet* genes; ■, strain B1 containing integrated *pet* genes; ○, M5A1 control. Figure 11B illustrates the production from cellobiose (100 g/liter) fermentation by strain P2. Symbols: ▲, ethanol; Δ, cell mass.

TABLE 17.

Ethanol production by recombinant strains of <i>K. oxytoca</i> M5A1 and by <i>E. coli</i> KO11.						
Organism ^a	Substrate	Additives ^b	Volumetric Productivity (g/liter/h)	Max. EtOH (g/liter)	Yield (g/g S)	% Theoretical ^c Yield
M5A1	10% glucose	none	0.72	16.2	0.17	33
M5A1 (pLOI555)	"	"	2.1	48.0	0.50	99
P2	"	"	1.6	46.4	0.46	90

^a Glucose and cellobiose fermentations were carried out at 30°C; other fermentations were carried out at 35°C.

^b Enzymes were added at the time of inoculation.

^c Computed for the time period between 6 and 24 hours.

^d Corrected for dilution by the addition of base during fermentation. S (substrate) refers to glucose, cellobiose or SOLKA FLOC SW40.

^e Based on a maximum theoretical yield of 51 g ethanol/100 g glucose, 53.5 g ethanol/100 g cellobiose, and 28 g ethanol/50 g cellulose. Values were not corrected for residual substrate.

^f SOLKA FLOC SW40 was predigested at 60°C for 12 h by using cells harvested from a prior fermentation as a source of *C. thermocellum* celD. After cooling to 35°C, broth was inoculated and 1 ml of commercial cellulase/100 ml fermentation volume added to start the fermentation.

TABLE 17. (continued)

Ethanol production by recombinant strains of <i>K. oxytoca</i> M5A1 and by <i>E. coli</i> KO11.						
Organism ^a	Substrate	Additives ^b	Volumetric Productivity (g/liter/h)	Max. EtOH (g/liter)	Yield (g/g S)	% Theoretical ^c Yield
B1	*	*	1.5	47.6	0.49	96
P2	10% cellobiose	*	1.5	45.2	0.49	96
B1	*	*	0.1	1.2	0.01	2
P2(pCT1057)	10% glucose	*	0.13	16.0	0.17	33
P2(pCT301)	*	*	0.58	19.4	0.20	40
P2(pCT603T)	*	*	0.83	30.0	0.32	62
P2	none	*	0.04	1.0	--	--
P2	none	10% CYTOLA SE	0.12	4.7	--	--
P2	none	10% MULTIP ECT	0.06	5.2	--	--
P2	5% SOLKA FLOC SW40	no enzyme	0.05	2.0	0.04	8
P2 (pCT603T) ^f	*	1.0% CYTOLA SE	0.09	17.4	0.36	65
P2 (pCT603T) ^f	*	1.0% MULTIF ECT	0.16	17.4	0.36	65
P2	5% SOLKA FLOC SW40	0.1% CYTOLA SE	0.08	2.3	0.07	12
P2	*	0.5% CYTOLA SE	0.13	7.0	0.14	26
P2	*	1.0% CYTOLA SE	0.18	8.7	0.18	32
P2	*	5.0% CYTOLA SE	0.52	16.9	0.36	64
P2	*	10% CYTOLA SE	0.50	16.3	0.35	62
P2	5% SOLKA FLOC SW40	0.1% MULTIF ECT	0.08	3.3	0.07	12
P2	*	0.5% Mutifect	0.18	5.9	0.13	23
P2	*	1.0% MULTIF ECT	0.36	10.4	0.23	41

^a Glucose and cellobiose fermentations were carried out at 30°C; other fermentations were carried out at 35°C.

^b Enzymes were added at the time of inoculation.

^c Computed for the time period between 6 and 24 hours.

^d Corrected for dilution by the addition of base during fermentation. S (substrate) refers to glucose, cellobiose or SOLKA FLOC SW40.

^e Based on a maximum theoretical yield of 51 g ethanol/100 g glucose, 53.5 g ethanol/100 g cellobiose, and 28 g ethanol/50 g cellulose. Values were not corrected for residual substrate.

^f SOLKA FLOC SW40 was predigested at 60°C for 12 h by using cells harvested from a prior fermentation as a source of *C. thermocellum* celD. After cooling to 35°C, broth was inoculated and 1 ml of commercial cellulase/100 ml fermentation volume added to start the fermentation.

TABLE 17. (continued)

Ethanol production by recombinant strains of <i>K. oxytoca</i> M5A1 and by <i>E. coli</i> KO11.						
Organism ^a	Substrate	Additives ^b	Volumetric Productivity (g/liter/h)	Max. EtOH (g/liter)	Yield (g/g S)	% Theoretical ^c Yield
P2	*	5.0% MULTIFECT	0.76	23.8	0.51	92
P2	*	10% MULTIFECT	0.86	32.5	0.69	123
KO11	5% SOLKA FLOC SW40	0.1% CYTOLASE	0.05	1.8	0.04	7
KO11	*	0.5% CYTOLASE	0.08	5.1	0.11	19
KO11	*	1.0% CYTOLASE	0.13	6.2	0.13	23
KO11	*	5.0% CYTOLASE	0.51	14.7	0.31	56
KO11	*	10% CYTOLASE	0.32	8.0	0.17	30
KO11	5% SOLKA FLOC SW40	0.1% MULTIFECT	0.06	2.2	0.05	8
KO11	*	0.5% MULTIFECT	0.08	2.7	0.06	10
KO11	*	1.0% MULTIFECT	0.18	5.6	0.12	21
KO11	*	5.0% MULTIFECT	0.43	20.6	0.43	76
KO11	*	10% MULTIFECT	0.84	30.7	0.63	112

^a Glucose and cellobiose fermentations were carried out at 30°C; other fermentations were carried out at 35°C.

^b Enzymes were added at the time of inoculation.

^c Computed for the time period between 6 and 24 hours.

^d Corrected for dilution by the addition of base during fermentation. S (substrate) refers to glucose, cellobiose or SOLKA FLOC SW40.

^e Based on a maximum theoretical yield of 51 g ethanol/100 g glucose, 53.5 g ethanol/100 g cellobiose, and 28 g ethanol/50 g cellulose. Values were not corrected for residual substrate.

^f SOLKA FLOC SW40 was predigested at 60°C for 12 h by using cells harvested from a prior fermentation as a source of *C. thermocellum* celLD. After cooling to 35°C, broth was inoculated and 1 ml of commercial cellulase/100 ml fermentation volume added to start the fermentation.

[0253] Comparison of crystalline cellulose fermentation (SOLKA FLOC SW40) by recombinant *K. oxytoca* M5A1 to *E. coli* in the presence of different concentrations of commercial cellulases: In contrast to ethanologenic *E. coli* which utilize only glucose, recombinant *K. oxytoca* P2 can ferment cellobiose and do not require exogenous β -glucosidase during cellulose fermentation. To examine the extent to which cellobiose utilization improves ethanol production from cellulose using commercial enzymes, fermentations were conducted with 50 g SOLKA FLOC SW40 as a substrate in the presence of either CYTOLASE or MULTIFECT concentrations from 0 to 10% (Fig. 12; Table 17). These preparations contain a mixture of fungal enzymes which include endoglucanases, exoglucanases, and β -glucosidases.

[0254] Figure 12 thus illustrates the effect of commercial cellulase additions on ethanol yield from 50 g SOLKA FLOC SW40/liter after 120 hours. The dotted line indicates the maximum amount of ethanol which can be made from only the added cellulose. The higher levels of ethanol are attributed primarily to additional substrate added as proprietary amendments in the MULTIFECT CL preparation. Symbols: \blacktriangle , fermented with *K. oxytoca* P2 and MULTIFECT; \blacksquare , fermented with strain *E. coli* KO11 and MULTIFECT; \triangle , fermented with *K. oxytoca* P2 and CYTOLASE; \square , fermented with *E. coli* KO11 and CYTOLASE.

[0255] At low concentrations, CYTOLASE appeared more efficient although some toxicity was associated with higher concentrations of this enzyme preparation. In all cases, the performance of the integrated strain of *K. oxytoca* was superior to *E. coli* KO11 despite the presence of β -glucosidase in commercial preparations of cellulase. Together, these results suggest that β -glucosidase activity in commercial cellulase preparations may not be saturating relative to the other cellulase activities.

[0256] With the highest concentration of MULTIFECT, yields exceeding the theoretical maximum from 50 g cellulose/l were observed. The source of this additional ethanol was investigated by conducting fermentations in Luria broth without added carbohydrate or enzymes and in Luria broth containing 10% CYTOLASE or 10% MULTIFECT but lacking added carbohydrate. Approximately 1 g ethanol/liter was produced from Luria broth alone and 5 g ethanol/liter from Luria broth containing 10% added cellulase. Thus with fermentations containing 10% MULTIFECT, approximately 4 g ethanol/liter can be attributed to the fermentation of commercial additives or enzyme stabilizing agents. It is likely that proportionate amounts of ethanol are also produced from lower concentrations of cellulases. Correcting for ethanol which could have been produced from Luria broth constituents and 10% MULTIFECT, ethanol yields from cellulose are approximately 94% of the theoretical maximum.

[0257] **Expression of *C. thermocellum* cellulases in strain P2:** Cellobiose can be fermented by strain P2 and is the dominant product from cellulose digestion by endoglucanases and exoglucanases. Although neither enzyme is made by our organism, integration of the pet genes into the chromosome facilitates the inclusion of plasmids encoding heterologous genes to provide these enzymes as co-products during fermentation. The expression of four endoglucanases from the thermophile, *Clostridium thermocellum*, was investigated after the addition of tetracycline resistance for selection (Table 18). The *celD* gene has been expressed at very high levels in *E. coli* and was expressed at high levels in strain P2. Most of the endoglucanase activity produced from these genes was retained within the cells although *Klebsiella* strains are known have protein secretion systems and the possibility of partial secretion was not eliminated.

[0258] A qualitative comparison of effectiveness in liquefaction of CMC was conducted in culture tubes during 48 hours incubation at 55°C. Consistent with activity measurements, recombinant cells harboring *celD* (pCT603T) were most effective followed by *celC* (pCT301). Strain P2 harboring *celB* (pCT207T) grew very poorly and was not tested further. *CelA* (pCT105T) recombinants caused little liquefaction.

[0259] The effects of producing heterologous gene products on fermentation performance were also investigated. Addition of plasmids expressing any one of the *cel* genes decreased the final cell density in 10% glucose fermentations by 10%-30% and decreased ethanol yield (Table 18). Expression of the *celD* gene was the least harmful with an ethanol yield of 0.32 g ethanol per gram of glucose, 62% of the theoretical maximum. Assays were conducted at 60°C using p-nitrophenyl- β -D-cellobioside as the substrate. Units are μ moles hydrolyzed per min per ml of fermentation broth containing cells or broth after cells had been removed by centrifugation.

TABLE 18.

Expression of <i>C. thermocellum</i> endoglucanases genes in strain P2.				
Gene	Plasmid	Activity (U/ml)		
		cells+broth	broth	% cell-associated
control	none	0.2	-	-
<i>celA</i>	pCT105T	2.3	0.4	83
<i>celB</i>	pCT207T	17	0.8	95
<i>celC</i>	pCT301	41	6.2	85
<i>celD</i>	pCT603T	49	13	73

[0260] **Hydrolysis and fermentation of amorphous cellulose by strain P2 expressing *celD*.** Although the *celD* gene product will not hydrolyze crystalline cellulose, this enzyme has been previously shown to hydrolyze amorphous cellulose (Bèguin *et al.*, Joliff *et al.*, *supra*). Strain P2(pCT603T) expressing this gene was tested for its ability to convert amorphous cellulose to ethanol. Phosphoric acid swollen and base swollen (sodium hydroxide) SOLKA FLOC SW40 were tested in a two-stage batch process using the cells of P2(pCT603T) from a prior glucose fermentation as a source of endoglucanase D and the same organism to produce ethanol.

[0261] The first step consisted of incubating 3 ml of Luria broth containing 76 mg/ml acid swollen cellulose or 32 mg/ml base swollen cellulose at 60°C (pH 6.0) for 72 hours using cells harvested from an equivalent volume from a prior glucose fermentation. Heating to 60°C inactivated cells to release endoglucanase D and provided a near optimal temperature for activity. The release of reducing sugars was measured during hydrolysis. Although substrate concentrations were different, acid swollen cellulose was hydrolyzed at twice the rate of base swollen cellulose to produce 240 μ molar and 60 μ molar cellobiose equivalents, respectively, after 24 hours. The acid swollen cellulose became quit

viscous making sampling difficult. After 72 hours, 110 μ molar cellobiose equivalents were present in the digestion of base swollen cellulose. These yields reflect the near complete hydrolysis of both acid swollen and base swollen cellulose.

[0262] Figures 13A, B and C comprise the thin layer chromatographic analyses of cellulose hydrolysis and fermentation by P2(pCT603T). Acid swollen cellulose (Fig. 13A), base swollen cellulose (Fig. 13B), and crystalline cellulose (Fig. 13C; SOLKA FLOC SW40) were tested. Cellobiose and glucose were used as standards in lane 1 of each chromatogram. In Figures 13A and B, lanes 2 through 6 represent samples removed at various times during hydrolysis of amorphous cellulose (0, 6, 12, 24, 48 and 72 hours, respectively); lane 7 represents broth analyzed in lane 6 after fermentation for 24 hours at 30°C with strain P2(pCT603T).

[0263] Figure 13C comprises the thin layer chromatographic analysis of crystalline cellulose (SOLKA FLOC SW40) fermentation with 1% MULTIFECT and P2(pCT603T) expressing *C. thermocellum celD*. The lane compositions are as follows: 2, SOLKA FLOC suspension immediately prior to digestion; 3, broth after 24 hours of digestion at 60°C with endoglucanase D supplied by P2(pCT603T) from a prior fermentation; 4, broth shown in lane 3 after fermentation for 24 hours at 35°C by P2(pCT603T); 5, after an additional 24 hours of fermentation by P2(pCT603T) at 35°C (total fermentation, 48 hours).

[0264] Thus, lanes 2-6 in figures 13A and 13B reveal the cellulose digestion products at various times from acid swollen and base swollen cellulose, respectively. Cellobiose is the dominant product in both with a small amount of glucose. Low levels of cellotriose also were present in digests of acid swollen cellulose. After cooling to 35°C, these broths were inoculated with 50 μ liter of P2(pCT603T) and allowed to ferment without agitation or pH control for 24 hours. Glucose, cellobiose, and cellotriose were removed completely during this period (Fig. 13A & B, lane 7) with the production of approximately 5 g ethanol/liter. For base swollen cellulose, 0.16 g ethanol/g cellulose was produced. Although the fall in pH probably served to limit the extent of ethanol production, this ethanol yield is equivalent to 27% of the theoretical maximum.

[0265] Partial replacement of commercial cellulase by endogenous production of *C. thermocellum* endoglucanase: Although P2(pCT603T) does not require externally added β -glucosidase and can provide endoglucanase D which will hydrolyze amorphous cellulose, additional activities are required to hydrolyze crystalline cellulose. The ability of this strain to partially replace the requirement for commercial cellulase was tested in a two stage fermentation process. Cells harvested from a prior glucose fermentation were resuspended in an equivalent volume of Luria broth containing 50 g SOLKA FLOC SW40/liter and incubated at 60°C for 24 hours. After cooling to 35°C, these fermentations were inoculated with the same organism along with 1% commercial cellulase. Similar results were obtained for both CYTOLASE and MULTIFECT with the production of 17.4 g ethanol/liter, 65% of the maximum theoretical yield (Table 17).

[0266] As illustrated in Figure 14 for CYTOLASE, pretreatment with the *celD* gene product greatly enhanced the ethanol production using 1% cellulase, providing an ethanol yield equivalent to that of 5% CYTOLASE. In Figure 14, cellulose (50 g SOLKA FLOC SW40/liter) was incompletely saccharified with CYTOLASE and endoglucanase D and fermented by strains P2 and P2(pCT603T). Symbols: Δ , 5 ml CYTOLASE added per 100 ml broth at time of inoculation and fermented with P2; \square , 1 ml CYTOLASE added per 100 ml broth at time of inoculation and fermented with P2; \circ , 0.1 ml CYTOLASE added per 100 ml broth at time of inoculation and fermented with P2; \blacksquare , SOLKA FLOC predigested at 60°C for 12 hours by using P2(pCT603T) cells harvested from a prior fermentation as a source of *C. thermocellum* endoglucanase D and fermented. After cooling to 35°C, 1 ml of commercial cellulase/100 ml fermentation broth and P2(pCT603T) were added to start the fermentation.

[0267] Figure 13C shows a thin layer chromatogram of samples from different stages of a parallel two step fermentation with MULTIFECT. Again, glucose, cellobiose, and cellotriose were completely metabolized. Thus the amount of commercial cellulase needed for saccharification was reduced by up to 80% after pretreatment with recombinant endoglucanase D produced as a co-product with ethanol in a prior fermentation.

[0268] Discussion: It is known that hydrolysis of cellobiose to monomeric sugar by β -glucosidase often limits cellulose digestion by fungal broths. The activity of β -glucosidase is typically the least abundant and least stable of the enzyme cadre required for efficient cellulose hydrolysis. The accumulation of cellobiose as a product of cellulase action acts as a competitive inhibitor of further depolymerization (Eriksson *et al.* [1990] Microbial and enzymatic degradation of wood and wood components. Springer-Verlag, New York; Jeffries, *supra*). *K. oxytoca* P2 is the first organism reported to rapidly and efficiently convert cellobiose to high levels of ethanol.

[0269] This organism appears to have the capacity to actively transport cellobiose and cellotriose, minimizing cellobiose accumulation and eliminating the requirement for further depolymerization by external enzymes. Although transport systems for cellobiose have not been investigated in *K. oxytoca* (Al [1989] *J. Biotechnol.* 12:79-86), the closely related organism *E. coli* contains genes which encode phosphotransferase components and phosphoglucosidase for cellobiose metabolism which are cryptic in most laboratory strains but function in nature (Hall *et al.* *J. Bacteriol.* 169: 2713-2717; Krickler *et al.* [1987] *Genetics* 115:419-429). It is likely that similar genes function in *K. oxytoca* (Al, *supra*).

[0270] Integration of the *pet* genes for ethanol production into the chromosome facilitated the production of plasmid-

borne, recombinant proteins as co-products with ethanol. Although cellulase genes from *C. thermocellum* were used as an example, other more valuable recombinant products could also be made such as lipases, proteinases, glycosidases, animal hormones, and biomedical products. The production of *C. thermocellum* cellulases by strain P2 as a co-product was accompanied by an unexpected decrease in the efficiency of fermentation. The basis for this decreased efficiency is unknown although the present inventors have made similar observations with ethanologenic *E. coli* containing the pullulanase gene from *Thermoanaerobium brockii* and with multi-plasmid ethanologenic constructs of *K. oxytoca* expressing the *C. thermocellum* xylanase gene.

[0271] The foregoing example with *K. oxytoca* thus forms the basis for an improved process to convert cellulose into ethanol. It is known that cellobiose and cellotriose are inhibitors of endoglucanase and exoglucanase, and that β -glucosidase is responsible for converting those oligomers into glucose monomers. Ethanologenic recombinants such as strain P2, however, do not require depolymerization of sugars for metabolism, eliminating the need for β -glucosidase and reducing end-product inhibition of cellulases by cellobiose and cellotriose. A further advantage is the reduction in potential contaminants since fewer organisms are able to metabolize cellobiose or cellotriose as compared to monomeric glucose. Based upon comparisons with *E. coli* KO11, which lack cellobiose utilization, the transport and metabolism of cellobiose exhibited by *K. oxytoca* P2 decreased the requirement for commercial enzymes during cellulose fermentation.

[0272] The requirement for commercial cellulases was further reduced by using supplemental endoglucanase produced as a co-product during fermentation. In this regard, thermostable enzymes appear particularly useful since they can be readily harvested within cells after fermentation and released in active form simply by heating to the temperature at which they function best. Pretreatment of SOLKA FLOC SW40 with endoglucanase D from a prior fermentation dramatically increased the effectiveness of Genencor cellulases. Since endoglucanase D is active only on amorphous regions of cellulose (Bèguin *et al.*, *supra*), the beneficial effect of this pretreatment may result from the creation of new sites for attack by commercial cellulases. In *C. thermocellum*, endoglucanase D is part of a cellulase complex (Bèguin *et al.*, *supra*). It is possible that this enzyme may also complex with fungal enzymes or bind cellulose and facilitate an opening of cellulose structure for hydrolysis.

[0273] The best construct for cellulose fermentation to ethanol, P2(pCT603T), is not as effective as native *C. thermocellum* (Tailliez *et al.* [1989] *Appl. Environ. Microbiol.* 55:203-211). Unlike *C. thermocellum*, *K. oxytoca* P2(pCT603T) lacks a complete cellulase system. However, in combination with 1% commercial CYTOLASE or MULTIFECT, P2 (pCT603) produced higher yields and final concentrations of ethanol at 35°C than the ethanol-tolerant *C. thermocellum* mutants at 65°C. Further genetic improvements or process improvements may be desirable.

Example 19 - Ethanol Production from Starch by Recombinant *Escherichia coli* containing Chromosomally Integrated *Z. mobilis* Genes for Ethanol Production and Plasmids Expressing Thermostable Genes for Saccharification

[0274] **Summary of Example:** Ethanologenic strains of *E. coli* have been developed which can express thermostable enzymes for starch saccharification as intracellular products. These enzymes can be harvested within cells at the end of fermentation and liberated by heating to the temperature at which they exhibit maximal activity (60°C to 70°C). Organisms such as these could be used to supply enzymes for yeast-based fermentations while producing a small amount of ethanol as a co-product.

[0275] In this example, advantage is taken of the absence of a protein secretion system in *E. coli* by developing ethanologenic strains which express genes encoding thermostable α -amylase and pullulanase as intracellular products. Thermostable enzymes are harvested within cells at the end of fermentation and are released for starch saccharification simply by heating to the temperature at which these enzymes exhibit near optimal activity.

[0276] **Organisms and growth conditions:** All ethanol fermentations were conducted in Luria broth with added carbohydrate using *E. coli* strain KO11 (Ohta *et al.*, 1991). Plasmids were added to this strain as indicated.

[0277] **Plasmid construction:** Two plasmids, pLOI140 and pLOI141 (Figs. 15A and B), were constructed which contained the α -amylase gene from *Bacillus stearothermophilus* (Mielenz *et al.* [1985] U.S. Patent No. 4,493,893) and the pullulanase gene from *Thermoanaerobium brockii* (Coleman *et al.* [1986] U.S. Patent No. 4,612,287; Coleman *et al.* [1987] *J. Bacteriol.* 169:4302-4307) using standard methods. A 5.4 kilobase pair (kbp) *Hind*III fragment from pWL625Amy (ATCC 31,791) containing the amylase gene was inserted into a *Hind*III site of pCPC902 (Coleman *et al.* [1986] *supra*) after partial digestion. Comparison of constructs on Luria agar containing starch and pullulan indicated that insertions at only one site allowed retention of high level expression for both genes, pLOI568. A 2.5 kbp *Eco*RI fragment from pCOS2EMBL (Poustka *et al.* [1984] *Proc. Natl. Acad. Sci. USA* 81:4129-4133) containing a tetracycline gene was added to pLOI568 after partial digestion with *Eco*RI to produce pLOI140 and pLOI141. The final plasmids contained a pBR322 replicon and were quite stable in strain KO11, being retained by 96% of the cells after 48 generations without antibiotic selection.

[0278] Figures 15A and B utilize the following abbreviations: amy, gene encoding *Bacillus stearothermophilus* α -amylase; pul, gene encoding *Thermoanaerobium brockii* pullulanase; tet, tetracycline resistance gene; amp, ampicillin

resistance gene; ori, *colE1* replicon.

[0279] **Fermentation and analyses:** Fermentations were carried out in 500-ml Fleakers which served as pH stats (350 ml working volume) essentially as described previously (Beall *et al.*, *supra*). Media consisted of Luria broth containing 10% glucose, 5% glucose, 10% maltose, 5% maltose, or 4.5% starch (30°C, pH 6.0, 100 rpm). Fermentations were inoculated to an initial cell density of 0.03 mg dry weight/ml. Sugars were sterilized separately by filtration. Starch was not sterilized.

[0280] Cell mass was measured as O.D. 550 nm (approximately 0.32 g cell dry weight/liter at 1 O.D.). Total saccharifying activity was measured at 60°C as the release of reducing sugars (Bergmeyer *et al.*, *supra*). Ethanol was determined by gas liquid chromatography (Dombek *et al.*, *supra*). Ethanol concentrations were expressed as g/liter. Ethanol yields were corrected for dilution by the addition of base during fermentations and were computed on the basis of total sugar or starch without correction for unused carbohydrate. Maximum theoretical yields from fermentation are approximately 0.51 g ethanol/g glucose, 0.53 g ethanol/g maltose, and 0.56 g ethanol/g starch. Results were based on an average of 2 or more fermentations.

[0281] **Fermentation of starch:** For starch fermentations, KO11(pLOI140) or KO11(pLOI141) cells were harvested from a prior 10% glucose fermentation after 72 hours and stored frozen. These cells were resuspended in 25 ml of Luria broth adjusted to pH 6.0. Starch (15.75 g) was suspended in 325 ml of Luria broth adjusted to pH 6.0. One-half of the cell suspension was added to the starch suspension, incubated in a boiling water bath with stirring until it reached 70°C, and held at 70°C for 5 minutes. This mixture was cooled to 60°C. Remaining cells were added and incubation continued for 24 hours at 60°C. Heating inactivated recombinant *E. coli* and released thermostable enzymes for starch hydrolysis. The saccharified starch was cooled to 30°C and adjusted to a volume of 350 ml with sterile distilled water.

[0282] An inoculum was grown overnight in 5% glucose, harvested by centrifugation, and used to start the fermentation (0.03 g cell dry weight of live cells/liter). Samples were removed for ethanol determination and for analysis by thin layer chromatography.

[0283] Glucose, maltose, maltotriose, and maltotetraose were separated by using unactivated, Whatman silica gel 150A plates (Whatman Inc., Clifton, New Jersey). Samples of 1 µliter containing 5 to 50 µg saccharide were applied and separated at room temperature by a single development in acetone, ethyl acetate, and acetic acid (2:1:1, respectively) to which 2% water was added immediately prior to use. Dried plates were stained as described by Bounias [1980] *Anal. Biochem.* 106:291-295. Starch and sugars were obtained from the Sigma Chemical Company, St. Louis, Mo.

[0284] **Fermentation of glucose and maltose by recombinant strains of *E. coli*:** The fermentation of glucose and maltose by KO11 was examined. The results are graphically illustrated in Figures 16A and B, which reflect ethanol production and cell mass, respectively. The symbols used in the figures are as follows: ●, strain KO11 with 10% glucose; Δ, strain KO11(pLOI140) with 10% glucose; □, strain KO11(pLOI141) with 10% glucose; ○, strain KO11 with 5% maltose (results with 5% and 10% maltose were identical).

[0285] Although the utilization of maltose by *E. coli* has been well-studied in aerobic cultures, growth and ethanol production from maltose were much slower than glucose reaching less than 25% of the maximal theoretical yield (53.5 g ethanol/liter) from 10% maltose and 50% of the maximal theoretical yield (26.8 g ethanol/liter) from 5% maltose. Since similar results were observed with both concentrations of maltose, it is unlikely that slower growth and fermentation result from direct maltose toxicity. The slower fermentation of maltose as compared to glucose may result from other limitations such as saccharide transport, glucose phosphorylation, etc.

[0286] As seen in Figures 16A and B, the addition of plasmids pLOI140 and pLOI141 encoding α-amylase and pullulanase to strain *E. coli* KO11 decreased the rates of growth and ethanol production from glucose. Final cell yields and ethanol yields after 72 hours were reduced by approximately 1/3.

[0287] **Expression of enzymes for saccharification:** Total saccharifying activities produced by KO11(pLOI140) and KO11(pLOI141) were measured at the end of fermentation (72 hours) as the release of reducing sugars using a suspension of potato starch in Luria broth (pH 6.0) as the substrate (70°C). Similar activity was observed with both strains, 30 µmoles reducing sugar/ml of culture per minute. Over 90% of this activity was intracellular but was readily released upon heating to 70°C.

[0288] **Fermentation of starch:** KO11(pLOI140) and KO11(pLOI141) were tested in a two-step process in which the enzymes for saccharification and ethanol production involved a single organism. Cells were harvested from a prior fermentation with 10% glucose, resuspended, and used as a source of enzymes to hydrolyze an equal volume of cold, 4.5% starch suspension (Luria broth, pH 6.0). Half of the harvested cells were added and the mixture rapidly heated to 70°C in a water bath for approximately 5 minutes. This mixture was shifted to 60°C and the balance of the cells added to facilitate hydrolysis. Corn, potato, rice, and wheat starch were tested and all behaved similarly. The starch suspension remained liquid throughout heating and hydrolysis in contrast to controls with KO11 lacking starch genes which solidified completely within 3 minutes of heating.

[0289] After incubation for 24 h at 60°C, the Luria broth containing hydrolyzed starch was cooled to 30°C and inoculated to start fermentation (pH 6.0, 100 rpm, inoculum of 0.03 g cell dry wt/liter) as previously described (Beall *et al.*,

1991, *supra*). Ethanol was produced at a similar rate from all types of starch, 0.26 to 0.28 g ethanol/liter during the initial 24 hours. Final ethanol yields were 0.35 to 0.37 g ethanol/gram of starch as compared to a theoretical maximum of 0.56 g ethanol/g starch. The results of the starch fermentation are provided in Figure 17.

[0290] Since the data for KO11(pLOI140) and KO11(pLOI141) were essentially identical, only the data for KO11 (pLOI141) is shown in Figure 17. The symbols used in Figure 17 are as follows: Δ , corn starch (S-4126); \square , potato starch (S-2004); \circ , rice starch (S-7260); \blacktriangle , wheat starch (S-5127); \blacksquare , corn starch with 2 mM CaCl_2 ; \bullet , KO11 with 5% glucose as a control.

[0291] Samples were removed at different stages of this two-step process with corn starch and analyzed by thin layer chromatography (Figure 18). After 24 hours, glucose, maltose, maltotriose, maltotetraose, and longer oligosaccharides are evident as hydrolysis products. Glucose, maltose, and maltotriose were utilized by KO11(pLOI141) during fermentation (72 hours).

[0292] Figure 18 comprises the thin layer chromatographic analysis of corn starch fermentation. Maltose and glucose served as standards. Glucose through maltotetraose are labeled G1 through G4, respectively. The lanes were as follows: 1, glucose and maltose as standards; 2, broth after 24 hours saccharification at 60°C; 3, broth shown in lane 2 after fermentation for 72 hours.

[0293] Starch-degrading enzymes are frequently stabilized by the presence of calcium ions. Fermentation experiments conducted with corn starch in Luria broth containing 2 mM calcium chloride resulted in slightly faster fermentation and a higher yield, 0.29 g ethanol/liter per hour and 0.40 g ethanol/g starch, respectively. As compared to conventional starch fermentations with yeasts and enzymes, the recombinant *E. coli* fermentation was slower and produced a lower final ethanol concentration. The overall ethanol yield from starch using this two-step process with a single organism was 85% that of current commercial processes using yeasts and supplemental enzymes (approximately 0.47 g ethanol/g starch).

[0294] **Conclusions:** While further improvements may be desirable, this example demonstrates that ethanologenic fermentations with recombinant *E. coli* could be used to produce part of the enzymes needed for starch saccharification. Enzyme production would be quite simple with no requirement for aeration. The thermostable bacterial enzymes function optimally at the same pH (approximately 6.0) as *E. coli*-based fermentation. With these organisms, residual starch in low value materials such as corn hulls could be used as a substrate and corn steep liquor as a source of complex nutrients. A process such as this may reduce the cost of enzymes for saccharification and increase ethanol yield from corn.

Claims

1. A recombinant host comprising heterologous DNA encoding:

- (a) alcohol dehydrogenase; and
- (b) pyruvate dehydrogenase; and
- (c) a polysaccharase or a combination of different polysaccharases;

wherein said host expresses said heterologous DNA at a sufficient level so as to facilitate the production of ethanol as a primary fermentation product by said host.

2. The recombinant host of claim 1 further comprising heterologous DNA encoding proteins involved in the transport of mono- and/or oligosaccharides into the host.

3. The recombinant host of claim 1 or claim 2 further comprising heterologous DNA encoding proteins which enable the host to transport and metabolize an oligosaccharide.

4. The host of any one of the preceding claims wherein said host is selected from bacteria, fungi and eukaryotic cells.

5. The host of claim 4 which is a bacterium selected from any of:

- (a) Gram-negative bacteria;
- (b) enteric bacteria;
- (c) *Erwinia* spp.;
- (d) *Klebsiella* spp.;
- (e) *Xanthomonas* spp..

6. The host of any one of the preceding claims wherein the polysaccharase(s) are selected from any of:

- (a) cellulolytic enzymes (e.g. endoglucanase, cellobiohydrolase or β -glucosidase);
- (b) hemicellulolytic enzymes;
- (c) xylanolytic enzymes (e.g. endo-1,4- β -xylanase, β -xylosidase, α -glucuronidase, α -L-arabinofuranosidase, acetylesterase or acetylxylanesterase);
- (d) starch-degrading enzymes (e.g. α -amylase (e.g. from *Bacillus stearothermophilus*), β -amylase, glucoamylase or pullulanase (e.g. from *Thermoanaerobium brockii*));
- (e) β -glucanase, arabinosidase, mannanase, pectin hydrolase or pectate lyase;
- (f) the expression product of a gene selected from *celD* (e.g. from *Clostridium thermocellum*), *xynZ* (e.g. from *Clostridium thermocellum*), *xyB* (e.g. from *Butyrivibrio fibrosolvens*) and a cellulase gene of *Cellulomonas fimi*.

7. The host of any one of the preceding claims wherein the polysaccharase(s) are:

- (a) thermostable, and/or
- (b) at least partially secreted from the host or substantially accumulated in said host.

8. The host of any one of the preceding claims wherein the heterologous DNA encoding the alcohol dehydrogenase and pyruvate dehydrogenase comprises the *Zymomonas mobilis* genes coding for alcohol dehydrogenase and pyruvate dehydrogenase.

9. A process for producing ethanol comprising fermentation with a culture of cells comprising the host of any one of the preceding claims such that ethanol is produced by the host as the primary fermentation product.

10. A plasmid comprising genes coding for:

- (a) alcohol dehydrogenase; and
- (b) pyruvate dehydrogenase; and
- (c) a polysaccharase or a combination of different polysaccharases;

wherein said plasmid is capable of directing a host to produce alcohol dehydrogenase, pyruvate dehydrogenase and the polysaccharase(s) at a sufficient level to facilitate the production of ethanol as the primary fermentation product by said host.

11. A method of treating an oligosaccharide feedstock, comprising the steps of:

- (a) providing a feedstock that contains a starting oligosaccharide;
- (b) bringing said feedstock into contact with an enzyme (e.g. a polysaccharase as defined in any one of claims 1-8) evolved by cells of a recombinant host according to any one of claims 1-8, such that said starting oligosaccharide is acted upon by said enzyme.

12. The method of claim 11 wherein said starting oligosaccharide is fermented to ethanol.

13. The method of claim 11 or 12 wherein said starting oligosaccharide:

- (a) comprises a disaccharide or trisaccharide;
- (b) is an insoluble oligosaccharide selected from cellulose, hemicellulose and starch.

14. The method of any one of claims 11-13 wherein step (b) comprises heating said cells to induce lysis and release of said enzyme (e.g. polysaccharase).

Patentansprüche

1. Rekombinanter Wirt mit heterologer DNA, die folgendes kodiert:

- (a) Alkoholdehydrogenase; und
- (b) Pyruvatdehydrogenase; und

(c) eine Polysaccharase oder eine Kombination unterschiedlicher Polysaccharasen;

wobei dieser Wirt die heterologe DNA auf einem ausreichenden Niveau exprimiert, um die Erzeugung von Ethanol als primärem Gärungsprodukt durch diesen Wirt zu erleichtern.

2. Rekombinanter Wirt nach Anspruch 1, der weiterhin heterologe DNA umfaßt, die Proteine kodiert, die am Transport von Mono- und/oder Oligosacchariden in den Wirt beteiligt sind.
3. Rekombinanter Wirt nach Anspruch 1 oder Anspruch 2, der weiterhin heterologe DNA umfaßt, die Proteine kodiert, die es dem Wirt ermöglichen, ein Oligosaccharid zu transportieren und metabolisieren.
4. Wirt nach einem der vorhergehenden Ansprüche, bei dem der Wirt aus Bakterien, Pilzen und eukaryotischen Zellen ausgewählt ist.
5. Wirt nach Anspruch 4, der ein aus folgendem ausgewähltes Bakterium ist:
 - (a) Gram-negatives Bakterien;
 - (b) Darmbakterien;
 - (c) *Erwinia* spp.;
 - (d) *Klebsiella* spp.;
 - (e) *Xanthomonas* spp.
6. Wirt nach einem der vorhergehenden Ansprüche, bei dem die Polysaccharase(n) aus folgendem ausgewählt ist:
 - (a) zellulolytischen Enzymen (beispielsweise Endoglucanase, Zellobiohydrolase oder β -Glucosidase);
 - (b) hemizellulolytischen Enzymen;
 - (c) xylanolytischen Enzymen (beispielsweise Endo-1,4- β -xylanase, β -Xylosidase, α -Glucuronidase, α -L-Arabinofuranosidase, Acetylsterase oder Acetylxylosterase);
 - (d) Stärke-abbauenden Enzymen (beispielsweise α -Amylase (beispielsweise von *Bacillus stearothermophilus*) β -Amylase, Glucoamylase oder Pullulanase (beispielsweise von *Thermoanaerobium brockii*);
 - (e) β -Glucanase, Arabinosidase, Mannanase, Pectinhydrolase oder Pectatlyase;
 - (f) dem Expressionsprodukt eines Gens, das aus *celD* (beispielsweise von *Clostridium thermocellum*), *xynZ* (beispielsweise von *Clostridium thermocellum*), *xylB* (beispielsweise von *Butyrivibrio fibrosolvens*) und einem Zellulase-Gen von *Cellulomonas fimi* ausgewählt ist.
7. Wirt nach einem der vorhergehenden Ansprüche, bei dem die Polysaccharase(n) folgendes sind:
 - (a) thermostabil und/oder
 - (b) zumindest teilweise vom Wirt sezerniert oder im wesentlichen in dem Wirt akkumuliert.
8. Wirt nach einem der vorhergehenden Ansprüche, bei dem die heterologe DNA, die die Alkoholdehydrogenase und Pyruvatdehydrogenase kodiert, die *Zymomonas mobilis*-Gene umfaßt, die Alkoholdehydrogenase und Pyruvatdehydrogenase kodieren.
9. Verfahren zur Erzeugung von Ethanol, das Gärung mit einer Kultur von Zellen umfaßt, die den Wirt nach einem der vorhergehenden Ansprüche umfassen, derart, das Ethanol durch den Wirt als das primäre Gärungsprodukt erzeugt wird.
10. Plasmid, das Gene umfaßt, die folgendes kodieren:
 - (a) Alkoholdehydrogenase; und
 - (b) Pyruvatdehydrogenase; und
 - (c) eine Polysaccharase oder eine Kombination unterschiedlicher Polysaccharasen;

wobei das Plasmid dazu in der Lage ist, einen Wirt zur Erzeugung von Ethanol durch Alkoholdehydrogenase, Pyruvatdehydro-

genase und der Polysaccharase(n) auf einem ausreichenden Niveau zu leiten, um die Erzeugung von Ethanol als dem primären Gärungsprodukt dieses Wirts zu erleichtern.

11. Verfahren zur Behandlung eines Oligosaccharid-Einsatzmaterials, das die folgenden Schritte umfaßt:

(a) Bereitstellen eines Einsatzmaterials, das ein Ausgangs-Oligosaccharid enthält;

(b) In Berührung bringen des Einsatzmaterials mit einem Enzym (beispielsweise einer Polysaccharase, wie sie in einem der Ansprüche 1 bis 8 definiert ist), das sich aus Zellen eines rekombinanten Wirts nach einem der Ansprüche 1 bis 8 entwickelt hat, derartig, daß das Ausgangs-Oligosaccharid von dem Enzym behandelt wird.

12. Verfahren nach Anspruch 11, bei dem das Ausgangs-Oligosaccharid zu Ethanol vergoren wird.

13. Verfahren nach Anspruch 11 oder 12, bei dem das Ausgangs-Oligosaccharid

(a) ein Disaccharid oder Trisaccharid umfaßt;

(b) ein unlösliches Oligosaccharid ist, das aus Zellulose, Hemizellulose und Stärke ausgewählt ist.

14. Verfahren nach einem der Ansprüche 11 bis 13, bei dem Schritt (b) ein Erhitzen der Zellen zur Induktion einer Lyse und Freisetzung des Enzyms (beispielsweise Polysaccharase) umfaßt.

Revendications

1. Un hôte recombinant comprenant du DNA hétérologue codant pour:

(a) Une alcool déshydrogénase; et

(b) Une pyruvate déshydrogénase; et

(c) Une polysaccharase ou une combinaison de différentes polysaccharases;

dans laquelle cet hôte exprime ce DNA hétérologue à un niveau suffisant de manière à faciliter la production d'éthanol par cet hôte sous forme d'un produit de fermentation primaire.

2. L'hôte recombinant selon la revendication 1 comprenant en outre un DNA hétérologue codant pour des protéines impliquées dans le transport de mono-et/ou oligosaccharides à l'intérieur de l'hôte.

3. L'hôte recombinant selon la revendication 1 ou 2 comprenant en outre du DNA hétérologue codant pour des protéines qui permettent à l'hôte de transporter et de métaboliser un oligosaccharide.

4. L'hôte selon l'une quelconque des revendications précédentes dans lequel cet hôte est choisi parmi les bactéries, les champignons et les cellules eucariotiques.

5. L'hôte selon la revendication 4 qui est une bactérie choisie parmi l'une quelconque des:

(a) Bactéries Gram-négatifs;

(b) entéobactéries;

(c) *Erwinia* spp.;

(d) *Klebsiella* spp.;

(e) *Xanthomonas* spp..

6. L'hôte selon l'une quelconque des revendications précédentes dans lequel les polysaccharases sont choisies parmi l'une quelconque des/de la/du:

(a) enzymes cellulolytiques (par exemple endoglucanase, cellobiohydrolase ou β -glucosidase);

(b) enzymes hémicellulolytiques;

(c) enzymes xylanolytiques (par exemple endo-1,4- β -xylanase, β -xylosidase, α -glucuronidase, α -L-arabinofuranosidase, acétyl estérase ou acétylxylanestérase);

(d) enzymes dégradant l'amidon (par exemple α -amylase (par exemple à partir de *Bacillus stearothermophilus*), β -amylase, glucoamylase ou pullulanase (par exemple à partir de *Thermoanaerobium brockii*));

(e) β -glucanase, arabinosidase, mannanase, pectinase ou pectate lyase;

(f) produit d'expression d'un gène choisi à partir de *celD* (par exemple à partir de *Clostridium thermocellum*), *xynZ* (par exemple à partir de *Clostridium thermocellum*), *xylB* (par exemple à partir de *Butyrivibrio fibrosolvens*) et un gène codant pour la cellulase de *Cellulomonas fimi*.

7. L'hôte selon l'une quelconque des revendications précédentes dans lequel les polysaccharases sont:

(a) thermostables, et/ou

(b) au moins partiellement sécrétées à partir de l'hôte ou essentiellement accumulées dans cet hôte.

8. L'hôte selon l'une quelconque des revendications précédentes dans lequel le DNA hétérologue codant pour l'alcool déshydrogénase et la pyruvate déshydrogénase comprend les gènes de *Zymomonas mobilis* codant pour l'alcool déshydrogénase et la pyruvate déshydrogénase.

9. Un procédé pour produire de l'éthanol comprenant la fermentation par une culture de cellules comprenant l'hôte de l'une quelconque des revendications précédentes telles que l'éthanol est produit par l'hôte sous forme de produit de fermentation primaire.

10. Un plasmide comprenant les gènes codant pour:

(a) l'alcool déshydrogénase; et

(b) la pyruvate déshydrogénase; et

(c) une polysaccharase ou une combinaison de différentes polysaccharases; dans laquelle ce plasmide est capable d'influencer un hôte à produire une alcool déshydrogénase, pyruvate déshydrogénase et la/les polysaccharase(s) à un niveau suffisant pour faciliter la production d'éthanol par cet hôte sous forme de produit de fermentation primaire.

11. Une méthode de traitement d'une charge d'un oligosaccharide, comprenant les étapes de:

(a) Fournir une charge qui contient un oligosaccharide de départ;

(b) mettre cette charge en contact avec une enzyme (par exemple une polysaccharase telle que définie dans l'une quelconque des revendications 1 à 8) produite par des cellules d'un hôte recombinant selon l'une quelconque des revendications 1 à 8, de sorte que cet oligosaccharide de départ soit traité par cette enzyme.

12. La méthode selon la revendication 11 dans laquelle cet oligosaccharide de départ est fermenté sous forme d'éthanol.

13. Méthode selon les revendications 11 ou 12 dans laquelle cet oligosaccharide de départ:

(a) comprend un disaccharide ou trisaccharide;

(b) est un oligosaccharide insoluble choisi parmi la cellulose, l'hémicellulose et l'amidon.

- 5 14. Méthode selon l'une quelconque des revendications 11 à 13 dans laquelle l'étape (b) comprend le chauffage de ces cellules pour induire la lyse et la libération de cette enzyme (par exemple la polysaccharase).

10

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C

SYMBOLS: ■, pLOI295; ●, pLOI308-2; ○, pLOI308-5,
▲, pLOI308-10; □, STRAIN TC4 LACKING A PLASMID




FIG. 4

SYMBOLS: ▲, AEROBIC GROWTH CONDITIONS;
●, ANAEROBIC GROWTH CONDITIONS

FIG. 5




FIG. 6





FIG. 7A




FIG. 7B



FIG. 7C





S 0 24 48 65
FIG. 8B

S 0 24 48 65
FIG. 8A




FIG. 10A




FIG. 10B




FIG. IIA





FIG. IIB






FIG. 12




FIG. 13C




FIG. 13B




FIG. 13A




FIG. 14

FIG. 15A




FIG. 15B




FIG. 16A





FIG. 16B






FIG. 17




FIG. 18

FIG. 19

